



EBP INTAKE ASSESSMENT PACKET MATCH-ADTC

Ages 5-6 Years

English

Required Forms
1. Demographic Information:
Client Intake Face Sheet □
2. Child's Top Problems:
Top Problems Assessment- Caregiver Report □
Top Problems Assessment – Child Report □
3. Child's Behavior & Functioning:
Ohio – Caregiver Report □
4. Child's Trauma History:
Trauma History Screen – Caregiver Report □
Supplemental Assessments
(Included in Packet)
Child Trauma Symptoms: YCPC- Caregiver Report
Child Dannessian, SMEO, Canagiyan Banant
Child Depression: <i>SMFQ</i> - Caregiver Report
Supplemental Assessments
(Included in Supplemental Assessment Packet)
Caregiver Symptoms:
PSS (Caregiver Stress)
PCL-5 (Caregiver Trauma Symptoms)
CESD-R (Caregiver Depression)



Intake Facesheet



VALIDATION REQUIREMENTS AND SYMBOLS EXPLAINED

- ! This symbol means the field is one of the minimum fields that must be filled out to save the record. No data will be saved unless these fields are completed.
- This symbol means the field is a required field in order to save the record as completed. Although you can save the record, the system does not consider the record to be completed unless ALL of these fields are completed.

		Direct Ser	vice Pr	ovide	r Use	r Information	1			
Clinician First and Last Name:										
Treatment Setting: Circle only ONE		strative Based School nity Support	CYFSC DCF Detention Extended	-		Group Home Hospital In-Home Outpatient Clinic	•	ntial T	entialTreatment Facility reatment Center d	Shelter Training Only Other
			Child	Infor	matic	on				
First Initial Child's First Name:				First I	Initial (Child's Last Nam	ie: <u>I</u>			
Date of Birth: !				Age:						
Sex: !	П	Female		_	Interse	х				
		Male			Other (specify)→				
Grade (current): *										
Race: *	_	American Indian Native	or Alaska	_	Black o	r African American			White	
		Asian		0		Hawaiian or Other Islander		_	Other (specify)	
Hispanic Origin: *		Yes, Cuban			Yes, of	Hispanic/Latino Orig	gin		Yes, South or Central Ar	merican
		Yes, Mexican, Me American, Chicar			Yes, Pu	erto Rican		_	No, Not of Hispanic, Lat Spanish Origin	ino, or
City/town:				ST:			Z *	ip:		
		Ch	ild Ide	ntifica	ation	Codes				
Agency-assigned Client ID Number (not PHI): !				PSDC	RS Clie	nt ID Number: !	!			
			Family	y Info	rmati	on				
Caregiver 1 Relationship: *				Careg	iver 2	Relationship:				
Preferred Language of Adult Participating in Treatment: *										
Does the adult participating in t	reatmo	ent speak Engli	ish?		Yes				No	
Primary Language of Child:							•	'		
Family Composition: * Select the choice that best describes		Two parent famil	у			parent - cal/adoptive parent			Relative/guardian	
the composition of the family.		Single Parent wit unrelated partne			Blende	d Family			Other	



Intake Facesheet



Living Situation of Child: *		College Dormitory		Job Corps		Psychiatric Hospital
What is the child's living situation?		Crisis Residence		Medical Hospital		Residential Treatment Facility
		DCF Foster Home		Mentor		TFC Foster Home (privately licensed)
		Group Home		Military Housing		Transitional Housing
		Homeless/Shelter		Other (specify):		
		Jail/Correctional Facility		Private Residence		
		System	Invo	olvement		
Child/Family involved with DCF?	*			Yes		No
If child / family is involved with	DCF, p	lease complete ALL of t	he fol	lowing questions:		
DCF Case ID: (if available)			_	Person Link ID: vailable)		
		Child Protective Services – In-Home		Family with Service Needs – (FWSN) In-Home		Not DCF – On Probation
DCF Status:	0	Child Protective Services – Out of Home		Family with Service Needs (FWSN) Out of Home	_	Not DCF – Other Court Involved
Dei Status.	0	Dual Commitment (JJ and Child Protective Services)		Juvenile Justice (delinquency) commitment	П	Termination of Parental Rights
		Family Assessment Response		Not DCF		Voluntary Services Program
DCF Regional Office:						
Youth involved with Juvenile Jus	stice (J	J) System? *		Yes		No
If youth is involved with JJ, pleas	se com	plete ALL of the followi	ing qu	estions:		
CSSD Client ID: (if available)			CSSE	Case ID: (if available)		
CSSD Case Type:				Delinquency		Family with Service Needs (Status Offense)
		Administrative Supervision		Juvenile probation	_	Restore Probation
CSSD Case Status:	0	Extended Probation		Non-Judicial FWSN Family Service Agreement	_	Suspended Order
coop case status.		Interim Orders		Non-Judicial Supervision (NJS)		Waived PDS - Probation
		Judicial FWSN Supervision		Non-Judicial Supervision Agreement		
Court District:						
Court Handling Decision:				Judicial	_	Non-Judicial
		Specific Trea	tmei	nt Information		
What treatment model are you	using v	with this child? *		TF-CBT		MATCH-ADTC
First Clinical Session Date: * Date of first EBP clinical session						



Intake Facesheet



		Treatme	nt In	formation	
Agency Referral Date/Request for Service: * Date child was referred to agency				icy Intake Date: * is the intake date for the client at ency?	
Referral Date: * Date referred for EBP services			Intal	ke Date: ! EBP Intake Date	
Referral Source: * Select the source of the EBP referral	П	Child Youth-Family Support Center (CYFSC)		Family Advocate	Physician
		Community Natural Support		Foster Parent	Police
		Congregate Care Facility		Info-Line (211)	Probation/Court
		CTBHP/Insurer		Juvenile Probation / Court	Psychiatric Hospital
		DCF	П	Other Community Provider Agency	School
		Detention Involved		Other Program within Agency	Self/Family
		Emergency Department		Other State Agency	
Assessment Outcome: What was the outcome of the referral to		Assessment not completed		Not appropriate for selected EBP	No treatment needed
the agency's EBP team? *	_	Appropriate for selected EBP		Not appropriate for selected EBP but needs other treatment	
CGI: Considering your experie of Intake? Circle only ONE:* Normal Slightly Severe Mile	nce, h			otional, behavioral, and/or farkedly Severe Very Sever	Among the most severe symptoms that any child may experience
		Treatment I	nforr	mation: School	
During the 3 months prior to the start of	EBP tre	eatment			
Child's school attendance: *	П	Good (few or no days missed)		No School Attendance: Child Too Young for School	No School Attendance: Other
		Fair (several days missed)		No School Attendance: Child Suspended/Expelled from School	
		Poor (many days missed)	0	No School Attendance: Child Dropped Out of School	
Suspended or expelled: *				Yes	No
IEP: *Does the child have an Individual	Educati	on Plan (special education)?		Yes	No
		Treatment I	nfor	mation: Legal	
During the 3 months prior to the start of	EBP tre			<u>_</u>	
Arrested: * Has the child been arrest	ed since	start of treatment?	_	Yes	No
Detained or incarcerated: * Has incarcerated since start of treatment?	the child	d been detained or		Yes	No
		Treatment In	form	ation: Medical	
During the 3 months prior to the start of	EBP tre	eatment			
Alcohol and/or drugs problems:	*			Yes	No
Evaluated in ER/ED for psychiati	ric issu	es: *	П	Yes	No
Certified medically complex: *	luated in ER/ED for psychiatric issues: *			Yes	No

Client ID:

Date of Completion: ___/___/___

Copyright © 2015, President and Fellows of Harvard College. All Rights Reserved.

Client Initials: _____

1

2

3

These copyrighted works were created by John R. Weisz, Ph.D., Kristel Thomassin, Ph.D., Jacqueline Hersh, Ph.D., and Rachel Vaughn-Coaxum, M.A., of Harvard's Laboratory for Youth Mental Health.

Client Initials:	Client ID:	Date of Completion://
Top Problems Assessment (TP	A) for MATCH-ADTC	
CHILD ASSESSMENT (English	sh)	

	Please enter each top problem in the text box below. How much have you had each of the following problems during the past week? Use a 0 to 4 scale. O=not a problem 4=a very big problem							
Rank	Top Problem	Rating (0-4)						
1								
2								
3								

Copyright © 2015, President and Fellows of Harvard College. All Rights Reserved.

These copyrighted works were created by John R. Weisz, Ph.D., Kristel Thomassin, Ph.D., Jacqueline Hersh, Ph.D., and Rachel Vaughn-Coaxum, M.A., of Harvard's Laboratory for Youth Mental Health.

Client Initials:	Client ID:	Date of Completion:	/	/

Ohio Mental Health Consumer Outcomes System Ohio Youth Problem and Functioning Scales (Caregiver: English) Parent Rating – Short Form

Instructions: Please rate the degree to which your child has experienced the following problems in the past 30 days.	Not at All	Once or Twice	Several Times	Often	Most of the Time	All of the Time
Arguing with others	0	1	2	3	4	5
2. Getting into fights	0	1	2	3	4	5
3. Yelling, swearing, or screaming at others	0	1	2	3	4	5
4. Fits of anger	0	1	2	3	4	5
5. Refusing to do things teachers or parents ask	0	1	2	3	4	5
6. Causing trouble for no reason	0	1	2	3	4	5
7. Using drugs or alcohol	0	1	2	3	4	5
8. Breaking rules or breaking the law (out past curfew, stealing)	0	1	2	3	4	5
9. Skipping school or classes	0	1	2	3	4	5
10. Lying	0	1	2	3	4	5
11. Can't seem to sit still, having too much energy	0	1	2	3	4	5
12. Hurting self (cutting or scratching self, taking pills)	0	1	2	3	4	5
13. Talking or thinking about death	0	1	2	3	4	5
14. Feeling worthless or useless	0	1	2	3	4	5
15. Feeling lonely and having no friends	0	1	2	3	4	5
16. Feeling anxious or fearful	0	1	2	3	4	5
17. Worrying that something bad is going to happen	0	1	2	3	4	5
18. Feeling sad or depressed	0	1	2	3	4	5
19. Nightmares	0	1	2	3	4	5
20. Eating problems	0	1	2	3	4	5

Copyright © Benjamin M. Ogles

(Add rating	gs together) To	otal
/	,	

January 2000 (Parent-1)

Response Scale for OHIO Problem Scale

0 1 2 3 4 5

Not at all Once or twice times Often Most of the time the time

Client Initials:	Client ID:	Date of Completion: / /	

Ohio Youth Problem and Functioning Scales (Caregiver: English)

Parent Rating – Short Form continued

Instructions: Please rate the degree to which your child's problems affect his or her current ability in everyday activities. Consider your child's current level of functioning.	Extreme Troubles	Quite a Few Troubles	Some Troubles	OK	Doing Very Well
Getting along with friends	0	1	2	3	4
2. Getting along with family	0	1	2	3	4
Dating or developing relationships with boyfriends orgirlfriends	0	1	2	3	4
Getting along with adults outside the family (teachers, principal)	0	1	2	3	4
5. Keeping neat and clean, looking good	0	1	2	3	4
6. Caring for health needs and keeping good health habits (taking medicines or brushing teeth)	0	1	2	3	4
7. Controlling emotions and staying out of trouble	0	1	2	3	4
Being motivated and finishing projects	0	1	2	3	4
Participating in hobbies (baseball cards, coins, stamps, art)	0	1	2	3	4
10. Participating in recreational activities (sports, swimming, bike riding)	0	1	2	3	4
11. Completing household chores (cleaning room, other chores)	0	1	2	3	4
12. Attending school and getting passing grades in school	0	1	2	3	4
13. Learning skills that will be useful for future jobs	0	1	2	3	4
14. Feeling good about self	0	1	2	3	4
15. Thinking clearly and making good decisions	0	1	2	3	4
16. Concentrating, paying attention, and completing tasks	0	1	2	3	4
17. Earning money and learning how to use money wisely	0	1	2	3	4
18. Doing things without supervision or restrictions	0	1	2	3	4
19. Accepting responsibility for actions	0	1	2	3	4
20. Ability to express feelings	0	1	2	3	4

Copyright	©	Benjamin	M.	Ogles
-----------	---	----------	----	-------

January 2000 (Parent-2)

(Add ratings together)	Total
------------------------	-------

Response Scale for OHIO Functioning Scale

0 1 2 3 4

Extreme Quite a few Some OK Doing troubles troubles troubles very well

			_	_
Client Initials:	Client ID:	Date of Completion:	/	/
Cilcrit illitials	CIICITE ID	Date of Completion.	_/	<i>'</i>

Trauma History Screen (THS) (Caregiver: English)

	Directions: Ask how many times each event happened, and how much it affected the child when it happened and now.	How many times has this happened?			The worst time this happened, how much did it affect him/her?					How much does this still affect your child?						
	"Has your child ever"	Never	Once	2-3 times	4-10 times	10+ times	Not at all	A little bit	Moderately	Quite a bit	Extremely	Not at all	A little bit	Moderately	Quite a bit	Extremely
1	Been in or seen a very bad accident?						1	2	3	4	5	1	2	3	4	5
2	Had someone s/he know been so badly injured or sick that s/he almost died?						1	2	3	4	5	1	2	3	4	5
3	Known somebody who died?						1	2	3	4	5	1	2	3	4	5
4	Been so sick or hurt that you or the doctor thought s/he might die?						1	2	3	4	5	1	2	3	4	5
5	Been unexpectedly separated from someone who s/he depends on for love or security for more than a few days?						1	2	3	4	5	1	2	3	4	5
6	Had somebody close to him/her try to kill or hurt themself?						1	2	3	4	5	1	2	3	4	5
7	Been physically hurt or threatened by someone?						1	2	3	4	5	1	2	3	4	5
8	Been robbed or seen someone get robbed?						1	2	3	4	5	1	2	3	4	5
9	Been kidnapped by somebody?						1	2	3	4	5	1	2	3	4	5
1 0	Been in or seen a hurricane, earthquake, tornado, or bad fire?						1	2	3	4	5	1	2	3	4	5
1	Been attacked by a dog or other animal?						1	2	3	4	5	1	2	3	4	5
1 2	Seen or heard people physically fighting or threatening to hurt each other?						1	2	3	4	5	1	2	3	4	5
1 3	Seen or heard somebody shooting a gun, using a knife, or using another weapon?						1	2	3	4	5	1	2	3	4	5
1 4	Seen a family member arrested or in jail?						1	2	3	4	5	1	2	3	4	5
1 5	Had a time in your life when s/he did not have the right care (e.g. food, clothing, a place to live)?						1	2	3	4	5	1	2	3	4	5
1 6	Been forced to see or do something sexual?						1	2	3	4	5	1	2	3	4	5
1 7	Seen or heard someone else being forced to do something sexual?						1	2	3	4	5	1	2	3	4	5
1 8	Watched people using drugs (like smoking, sniffing, or using needles)?						1	2	3	4	5	1	2	3	4	5
1 9	Seen something else that was very scary or where s/he thought somebody might get hurt or die? Specify:						1	2	3	4	5	1	2	3	4	5

20. Which one **bothers your child the MOST** right now: #_____ How long ago did it happen: _____

Response Scale for THS

1 2 3 4 5

Not at All Bit Moderately Quite Extremely A bit

Client Initials:	Client ID:	Date of Completion: /	/

YOUNG CHILD PTSD CHECKLIST (YCPC) (Caregiver: English)

For Child under 7 years old.

Below is a list of symptoms that children can have after life-threatening events.

When you think of ALL the life-threatening traumatic events from the first page, circle the number below (0-4) that best describes how often the symptom has bothered you in the LAST 2 WEEKS.

Not at all Once in a while 2 to 4 times a week/ Malmost always Everyday		0	1	2	3			4		
1. Does your child have intrusive memories of the trauma? Does s/he bring it up on his/her own? 0 1 2 3 4 2. Does your child re-enact the trauma in play with dolls or toys? This would be seenes that look just like the trauma. Or does s/he act it out by him/herself or with other kids? 0 1 2 3 4 3. Is your child having more nightmares since the trauma(s) occurred? 0 1 2 3 4 4. Does your child act like the traumatic event is happening to him/her again, even when it isn't? This is where a child is acting like they are back in the traumatic event and aren't in touch with reality. This is a pretty obvious thing when it happens. 0 1 2 3 4 5. Since the trauma(s) has s/he had episodes when s/he seems to freeze? You may have tried to snap him/her out of it but s/he was unresponsive. 0 1 2 3 4 6. Does s/he get upset when exposed to reminders of the event(s)? For example, a child who was in a car wreck might be nervous when it is raining. Or, a child who saw domestic violence might be nervous when it is raining. Or, a child who saw domestic violence might be nervous when other people argue. Or, a girl who was sexually abused might be nervous when other people alk about what happened, does s/he walk away or change the topic? 0 1 2 3 4 8. Does your c		Not at all		2 to 4 times a week/		eek/		Ever	yday	
Up on his/her own? 2. Does your child re-enact the trauma in play with dolls or toys? This would be scenes that look just like the trauma. Or does s/he act it out by him/herself or with other kids? 3. Is your child having more nightmares since the trauma(s) occurred? 0			Once in a while	Half the time	Almost always					
Up on his/her own? 2. Does your child re-enact the trauma in play with dolls or toys? This would be scenes that look just like the trauma. Or does s/he act it out by him/herself or with other kids? 3. Is your child having more nightmares since the trauma(s) occurred? 0		T =						1 _	1 -	
Does your child re-enact the trauma in play with dolls or toys? This would be scenes that look just like the trauma. Or does s/he act it out by him/herself or with other kids? 3. Is your child having more nightmares since the trauma(s) occurred? 0 1 2 3 4	1.	_		nories of the trauma?	Does s/he bring it	0	1	2	3	4
be scenes that look just like the trauma. Or does s/he act it out by him/herself or with other kids? 3. Is your child having more nightmares since the trauma(s) occurred? 4. Does your child act like the traumatic event is happening to him/her again, even when it isn't? This is where a child is acting like they are back in the traumatic event and aren't in touch with reality. This is a pretty obvious thing when it happens. 5. Since the trauma(s) has s/he had episodes when s/he seems to freeze? You may have tried to snap him/her out of it but s/he was unresponsive. 6. Does s/he get upset when exposed to reminders of the event(s)? For example, a child who was in a hurricane might be nervous when it is raining. Or, a child who was in a hurricane might be nervous when other people argue. Or, a girl who was sexually abused might be nervous when someone touches her. 7. Does your child get physically distressed when exposed to reminders? Like heart racing, shaking hands, sweaty, short of breath, or sick to his/her stomach? Think of the same type of examples as in #6. 8. Does your child try to avoid conversations that might remind him/her of the trauma(s)? For example, if other people talk about what happened, does s/he walk away or change the topic? 9. Does your child try to avoid things or places that remind him/her of the trauma(s)? For example, a child who was in a car wreck might try to avoid getting into a car. Or, a child who was in a flood might tell you not to drive over a bridge. Or, a child who was in a flood might tell you not to drive over a bridge. Or, a child who was in a flood might tell you not to drive over a bridge. Or, a child who saw domestic violence might be nervous to go in the house where it occurred. Or, a girl who was sexually abused might be nervous about going to bed because that's where she was abused before. 10. Does your child have difficulty remembering the whole incident? Has s/he blocked out the entire event? 11. Has s/he lost interest in doing things that s/he used to like to		1			0.5511					
him/herself or with other kids? Is your child having more nightmares since the trauma(s) occurred? 0	2.	-			=	0	1	2	3	4
 3. Is your child having more nightmares since the trauma(s) occurred? 4. Does your child act like the traumatic event is happening to him/her again, even when it isn't? This is where a child is acting like they are back in the traumatic event and aren't in touch with reality. This is a pretty obvious thing when it happens. 5. Since the trauma(s) has s/he had episodes when s/he seems to freeze? You may have tried to snap him/her out of it but s/he was unresponsive. 6. Does s/he get upset when exposed to reminders of the event(s)? For example, a child who was in a car wreck might be nervous while riding in a car now. Or, a child who saw domestic violence might be nervous when other people argue. Or, a girl who was sexually abused might be nervous when other people argue. Or, a girl who was sexually abused might be nervous when other people argue. Or, a child who saw domestic violence might be nervous when other people argue. Or, a girl who was sexually abused might be nervous when other people argue. Or, a child who saw domestic violence might be nervous when other people argue. Or, a child voice of breath, or sick to his/her stomach? Think of the same type of examples as in #6. 8. Does your child try to avoid conversations that might remind him/her of the trauma(s)? For example, if other people talk about what happened, does s/he walk away or change the topic? 9. Does your child try to avoid things or places that remind him/her of the trauma(s)? For example, a child who was in a car wreck might try to avoid getting into a car. Or, a child who was in a flood might tell you not to drive over a bridge. Or, a child who saw domestic violence might be nervous to go in the house where it occurred. Or, a girl who was sexually abused might be nervous about going to bed because that's where she was abused before. 10. Does your child have difficulty remembering the whole incident? Has s/he blocked out the entire event? 11. Has s/he lost interest in doing thing				ıma. Or does s/he act	it out by					
 4. Does your child act like the traumatic event is happening to him/her again, even when it isn't? This is where a child is acting like they are back in the traumatic event and aren't in touch with reality. This is a pretty obvious thing when it happens. 5. Since the trauma(s) has s/he had episodes when s/he seems to freeze? You may have tried to snap him/her out of it but s/he was unresponsive. 6. Does s/he get upset when exposed to reminders of the event(s)? For example, a child who was in a car wreck might be nervous while riding in a car now. Or, a child who was in a hurricane might be nervous when other people argue. Or, a girl who was sexually abused might be nervous when someone touches her. 7. Does your child get physically distressed when exposed to reminders? Like heart racing, shaking hands, sweaty, short of breath, or sick to his/her stomach? Think of the same type of examples as in #6. 8. Does your child try to avoid conversations that might remind him/her of the trauma(s)? For example, if other people talk about what happened, does s/he walk away or change the topic? 9. Does your child try to avoid things or places that remind him/her of the trauma(s)? For example, a child who was in a flood might tell you not to drive over a bridge. Or, a child who saw domestic violence might be nervous to go in the house where it occurred. Or, a girl who was sexually abused might be nervous about going to bed because that's where she was abused before. 10. Does your child have difficulty remembering the whole incident? Has s/he blocked out the entire event? 11. Has s/he lost interest in doing things that s/he used to like to do since the trauma(s)? 12. Since the trauma(s), does your child show a restricted range of positive 0 1 2 3 4 										
even when it isn't? This is where a child is acting like they are back in the traumatic event and aren't in touch with reality. This is a pretty obvious thing when it happens. 5. Since the trauma(s) has s/he had episodes when s/he seems to freeze? You may have tried to snap him/her out of it but s/he was unresponsive. 6. Does s/he get upset when exposed to reminders of the event(s)? For example, a child who was in a car wreck might be nervous while riding in a car now. Or, a child who saw domestic violence might be nervous when other people argue. Or, a girl who was sexually abused might be nervous when someone touches her. 7. Does your child get physically distressed when exposed to reminders? Like heart racing, shaking hands, sweaty, short of breath, or sick to his/her stomach? Think of the same type of examples as in #6. 8. Does your child try to avoid conversations that might remind him/her of the trauma(s)? For example, if other people talk about what happened, does s/he walk away or change the topic? 9. Does your child try to avoid things or places that remind him/her of the trauma(s)? For example, a child who was in a car wreck might try to avoid getting into a car. Or, a child who was in a flood might tell you not to drive over a bridge. Or, a child who was man a flood might tell you not to drive over a bridge. Or, a child who saw domestic violence might be nervous to go in the house where it occurred. Or, a girl who was sexually abused might be nervous about going to bed because that's where she was abused before. 10. Does your child have difficulty remembering the whole incident? Has s/he blocked out the entire event? 11. Has s/he lost interest in doing things that s/he used to like to do since the trauma(s)? 12. Since the trauma(s), does your child show a restricted range of positive 0 1 2 3 4							1			
traumatic event and aren't in touch with reality. This is a pretty obvious thing when it happens. 5. Since the trauma(s) has s/he had episodes when s/he seems to freeze? You may have tried to snap him/her out of it but s/he was unresponsive. 6. Does s/he get upset when exposed to reminders of the event(s)? For example, a child who was in a car wreck might be nervous while riding in a car now. Or, a child who saw domestic violence might be nervous when other people argue. Or, a girl who was sexually abused might be nervous when someone touches her. 7. Does your child get physically distressed when exposed to reminders? Like heart racing, shaking hands, sweaty, short of breath, or sick to his/her stomach? Think of the same type of examples as in #6. 8. Does your child try to avoid conversations that might remind him/her of the trauma(s)? For example, if other people talk about what happened, does s/he walk away or change the topic? 9. Does your child try to avoid things or places that remind him/her of the trauma(s)? For example, a child who was in a car wreck might try to avoid getting into a car. Or, a child who was in a flood might tell you not to drive over a bridge. Or, a child who saw domestic violence might be nervous to go in the house where it occurred. Or, a girl who was sexually abused might be nervous about going to bed because that's where she was abused before. 10. Does your child have difficulty remembering the whole incident? Has s/he blocked out the entire event? 11. Has s/he lost interest in doing things that s/he used to like to do since the trauma(s)? 12. Since the trauma(s), does your child show a restricted range of positive 0 1 2 3 4	4.					0	1	2	3	4
thing when it happens. 5. Since the trauma(s) has s/he had episodes when s/he seems to freeze? You may have tried to snap him/her out of it but s/he was unresponsive. 6. Does s/he get upset when exposed to reminders of the event(s)? For example, a child who was in a car wreck might be nervous while riding in a car now. Or, a child who was in a hurricane might be nervous when it is raining. Or, a child who saw domestic violence might be nervous when other people argue. Or, a girl who was sexually abused might be nervous when someone touches her. 7. Does your child get physically distressed when exposed to reminders? Like heart racing, shaking hands, sweaty, short of breath, or sick to his/her stomach? Think of the same type of examples as in #6. 8. Does your child try to avoid conversations that might remind him/her of the trauma(s)? For example, if other people talk about what happened, does s/he walk away or change the topic? 9. Does your child try to avoid things or places that remind him/her of the walk away or change the topic? 9. Does your child try to avoid things or places that remind him/her of the trauma(s)? For example, a child who was in a car wreck might try to avoid getting into a car. Or, a child who was in a flood might tell you not to drive over a bridge. Or, a child who was domestic violence might be nervous to go in the house where it occurred. Or, a girl who was sexually abused might be nervous about going to bed because that's where she was abused before. 10. Does your child have difficulty remembering the whole incident? Has s/he blocked out the entire event? 11. Has s/he lost interest in doing things that s/he used to like to do since the trauma(s)? 12. Since the trauma(s), does your child show a restricted range of positive 0 1 2 3 4					-					
5. Since the trauma(s) has s/he had episodes when s/he seems to freeze? You may have tried to snap him/her out of it but s/he was unresponsive. 6. Does s/he get upset when exposed to reminders of the event(s)? For example, a child who was in a car wreck might be nervous while riding in a car now. Or, a child who was in a hurricane might be nervous when it is raining. Or, a child who saw domestic violence might be nervous when other people argue. Or, a girl who was sexually abused might be nervous when someone touches her. 7. Does your child get physically distressed when exposed to reminders? Like heart racing, shaking hands, sweaty, short of breath, or sick to his/her stomach? Think of the same type of examples as in #6. 8. Does your child try to avoid conversations that might remind him/her of the trauma(s)? For example, if other people talk about what happened, does s/he walk away or change the topic? 9. Does your child try to avoid things or places that remind him/her of the value and thing into a car. Or, a child who was in a car wreck might try to avoid getting into a car. Or, a child who was in a flood might tell you not to drive over a bridge. Or, a child who saw domestic violence might be nervous to go in the house where it occurred. Or, a girl who was sexually abused might be nervous about going to bed because that's where she was abused before. 10. Does your child have difficulty remembering the whole incident? Has s/he blocked out the entire event? 11. Has s/he lost interest in doing things that s/he used to like to do since the trauma(s)? 12. Since the trauma(s), does your child show a restricted range of positive 0 1 2 3 4				with reality. This is a	pretty obvious					
may have tried to snap him/her out of it but s/he was unresponsive. 6. Does s/he get upset when exposed to reminders of the event(s)? For example, a child who was in a car wreck might be nervous while riding in a car now. Or, a child who was in a hurricane might be nervous when it is raining. Or, a child who saw domestic violence might be nervous when other people argue. Or, a girl who was sexually abused might be nervous when someone touches her. 7. Does your child get physically distressed when exposed to reminders? Like heart racing, shaking hands, sweaty, short of breath, or sick to his/her stomach? Think of the same type of examples as in #6. 8. Does your child try to avoid conversations that might remind him/her of the trauma(s)? For example, if other people talk about what happened, does s/he walk away or change the topic? 9. Does your child try to avoid things or places that remind him/her of the trauma(s)? For example, a child who was in a car wreck might try to avoid getting into a car. Or, a child who was in a flood might tell you not to drive over a bridge. Or, a child who saw domestic violence might be nervous to go in the house where it occurred. Or, a girl who was sexually abused might be nervous about going to bed because that's where she was abused before. 10. Does your child have difficulty remembering the whole incident? Has s/he blocked out the entire event? 11. Has s/he lost interest in doing things that s/he used to like to do since the trauma(s)? 12. Since the trauma(s), does your child show a restricted range of positive 0 1 2 3 4										
6. Does s/he get upset when exposed to reminders of the event(s)? For example, a child who was in a car wreck might be nervous while riding in a car now. Or, a child who saw domestic violence might be nervous when other people argue. Or, a girl who was sexually abused might be nervous when someone touches her. 7. Does your child get physically distressed when exposed to reminders? Like heart racing, shaking hands, sweaty, short of breath, or sick to his/her stomach? Think of the same type of examples as in #6. 8. Does your child try to avoid conversations that might remind him/her of the trauma(s)? For example, if other people talk about what happened, does s/he walk away or change the topic? 9. Does your child try to avoid things or places that remind him/her of the trauma(s)? For example, a child who was in a car wreck might try to avoid getting into a car. Or, a child who was in a flood might tell you not to drive over a bridge. Or, a child who saw domestic violence might be nervous to go in the house where it occurred. Or, a girl who was sexually abused might be nervous about going to bed because that's where she was abused before. 10. Does your child have difficulty remembering the whole incident? Has s/he blocked out the entire event? 11. Has s/he lost interest in doing things that s/he used to like to do since the trauma(s)? 12. Since the trauma(s), does your child show a restricted range of positive 0 1 2 3 4	5.					0	1	2	3	4
example, a child who was in a car wreck might be nervous while riding in a car now. Or, a child who was in a hurricane might be nervous when it is raining. Or, a child who saw domestic violence might be nervous when other people argue. Or, a girl who was sexually abused might be nervous when someone touches her. 7. Does your child get physically distressed when exposed to reminders? Like heart racing, shaking hands, sweaty, short of breath, or sick to his/her stomach? Think of the same type of examples as in #6. 8. Does your child try to avoid conversations that might remind him/her of the trauma(s)? For example, if other people talk about what happened, does s/he walk away or change the topic? 9. Does your child try to avoid things or places that remind him/her of the trauma(s)? For example, a child who was in a car wreck might try to avoid getting into a car. Or, a child who was in a flood might tell you not to drive over a bridge. Or, a child who saw domestic violence might be nervous to go in the house where it occurred. Or, a girl who was sexually abused might be nervous about going to bed because that's where she was abused before. 10. Does your child have difficulty remembering the whole incident? Has s/he blocked out the entire event? 11. Has s/he lost interest in doing things that s/he used to like to do since the trauma(s)?										
car now. Or, a child who was in a hurricane might be nervous when it is raining. Or, a child who saw domestic violence might be nervous when other people argue. Or, a girl who was sexually abused might be nervous when someone touches her. 7. Does your child get physically distressed when exposed to reminders? Like heart racing, shaking hands, sweaty, short of breath, or sick to his/her stomach? Think of the same type of examples as in #6. 8. Does your child try to avoid conversations that might remind him/her of the trauma(s)? For example, if other people talk about what happened, does s/he walk away or change the topic? 9. Does your child try to avoid things or places that remind him/her of the trauma(s)? For example, a child who was in a car wreck might try to avoid getting into a car. Or, a child who was in a flood might tell you not to drive over a bridge. Or, a child who saw domestic violence might be nervous to go in the house where it occurred. Or, a girl who was sexually abused might be nervous about going to bed because that's where she was abused before. 10. Does your child have difficulty remembering the whole incident? Has s/he blocked out the entire event? 11. Has s/he lost interest in doing things that s/he used to like to do since the trauma(s)? 12. Since the trauma(s), does your child show a restricted range of positive 0 1 2 3 4	6.					0	1	2	3	4
raining. Or, a child who saw domestic violence might be nervous when other people argue. Or, a girl who was sexually abused might be nervous when someone touches her. 7. Does your child get physically distressed when exposed to reminders? Like heart racing, shaking hands, sweaty, short of breath, or sick to his/her stomach? Think of the same type of examples as in #6. 8. Does your child try to avoid conversations that might remind him/her of the trauma(s)? For example, if other people talk about what happened, does s/he walk away or change the topic? 9. Does your child try to avoid things or places that remind him/her of the trauma(s)? For example, a child who was in a car wreck might try to avoid getting into a car. Or, a child who was in a flood might tell you not to drive over a bridge. Or, a child who saw domestic violence might be nervous to go in the house where it occurred. Or, a girl who was sexually abused might be nervous about going to bed because that's where she was abused before. 10. Does your child have difficulty remembering the whole incident? Has s/he blocked out the entire event? 11. Has s/he lost interest in doing things that s/he used to like to do since the trauma(s)? 12. Since the trauma(s), does your child show a restricted range of positive 0 1 2 3 4		example, a ch	ild who was in a car v	wreck might be nervou	is while riding in a					
other people argue. Or, a girl who was sexually abused might be nervous when someone touches her. 7. Does your child get physically distressed when exposed to reminders? Like heart racing, shaking hands, sweaty, short of breath, or sick to his/her stomach? Think of the same type of examples as in #6. 8. Does your child try to avoid conversations that might remind him/her of the trauma(s)? For example, if other people talk about what happened, does s/he walk away or change the topic? 9. Does your child try to avoid things or places that remind him/her of the trauma(s)? For example, a child who was in a car wreck might try to avoid getting into a car. Or, a child who was in a flood might tell you not to drive over a bridge. Or, a child who saw domestic violence might be nervous to go in the house where it occurred. Or, a girl who was sexually abused might be nervous about going to bed because that's where she was abused before. 10. Does your child have difficulty remembering the whole incident? Has s/he blocked out the entire event? 11. Has s/he lost interest in doing things that s/he used to like to do since the trauma(s)? 12. Since the trauma(s), does your child show a restricted range of positive 0 1 2 3 4		car now. Or, a child who was in a hurricane might be nervous when it is								
when someone touches her. 7. Does your child get physically distressed when exposed to reminders? Like heart racing, shaking hands, sweaty, short of breath, or sick to his/her stomach? Think of the same type of examples as in #6. 8. Does your child try to avoid conversations that might remind him/her of the trauma(s)? For example, if other people talk about what happened, does s/he walk away or change the topic? 9. Does your child try to avoid things or places that remind him/her of the trauma(s)? For example, a child who was in a car wreck might try to avoid getting into a car. Or, a child who was in a flood might tell you not to drive over a bridge. Or, a child who saw domestic violence might be nervous to go in the house where it occurred. Or, a girl who was sexually abused might be nervous about going to bed because that's where she was abused before. 10. Does your child have difficulty remembering the whole incident? Has s/he blocked out the entire event? 11. Has s/he lost interest in doing things that s/he used to like to do since the trauma(s)? 12. Since the trauma(s), does your child show a restricted range of positive 0 1 2 3 4		raining. Or, a child who saw domestic violence might be nervous when								
 7. Does your child get physically distressed when exposed to reminders? Like heart racing, shaking hands, sweaty, short of breath, or sick to his/her stomach? Think of the same type of examples as in #6. 8. Does your child try to avoid conversations that might remind him/her of the trauma(s)? For example, if other people talk about what happened, does s/he walk away or change the topic? 9. Does your child try to avoid things or places that remind him/her of the trauma(s)? For example, a child who was in a car wreck might try to avoid getting into a car. Or, a child who was in a flood might tell you not to drive over a bridge. Or, a child who saw domestic violence might be nervous to go in the house where it occurred. Or, a girl who was sexually abused might be nervous about going to bed because that's where she was abused before. 10. Does your child have difficulty remembering the whole incident? Has s/he blocked out the entire event? 11. Has s/he lost interest in doing things that s/he used to like to do since the trauma(s)? 12. Since the trauma(s), does your child show a restricted range of positive 0 1 2 3 4 				vas sexually abused m	night be nervous					
heart racing, shaking hands, sweaty, short of breath, or sick to his/her stomach? Think of the same type of examples as in #6. 8. Does your child try to avoid conversations that might remind him/her of the trauma(s)? For example, if other people talk about what happened, does s/he walk away or change the topic? 9. Does your child try to avoid things or places that remind him/her of the trauma(s)? For example, a child who was in a car wreck might try to avoid getting into a car. Or, a child who was in a flood might tell you not to drive over a bridge. Or, a child who saw domestic violence might be nervous to go in the house where it occurred. Or, a girl who was sexually abused might be nervous about going to bed because that's where she was abused before. 10. Does your child have difficulty remembering the whole incident? Has s/he blocked out the entire event? 11. Has s/he lost interest in doing things that s/he used to like to do since the trauma(s)? 12. Since the trauma(s), does your child show a restricted range of positive 0 1 2 3 4		when someon	e touches her.							
stomach? Think of the same type of examples as in #6. 8. Does your child try to avoid conversations that might remind him/her of the trauma(s)? For example, if other people talk about what happened, does s/he walk away or change the topic? 9. Does your child try to avoid things or places that remind him/her of the trauma(s)? For example, a child who was in a car wreck might try to avoid getting into a car. Or, a child who was in a flood might tell you not to drive over a bridge. Or, a child who saw domestic violence might be nervous to go in the house where it occurred. Or, a girl who was sexually abused might be nervous about going to bed because that's where she was abused before. 10. Does your child have difficulty remembering the whole incident? Has s/he blocked out the entire event? 11. Has s/he lost interest in doing things that s/he used to like to do since the trauma(s)? 12. Since the trauma(s), does your child show a restricted range of positive 0 1 2 3 4	7.	Does your chi	ild get physically distr	ressed when exposed t	o reminders? Like	0	1	2	3	4
8. Does your child try to avoid conversations that might remind him/her of the trauma(s)? For example, if other people talk about what happened, does s/he walk away or change the topic? 9. Does your child try to avoid things or places that remind him/her of the trauma(s)? For example, a child who was in a car wreck might try to avoid getting into a car. Or, a child who was in a flood might tell you not to drive over a bridge. Or, a child who saw domestic violence might be nervous to go in the house where it occurred. Or, a girl who was sexually abused might be nervous about going to bed because that's where she was abused before. 10. Does your child have difficulty remembering the whole incident? Has s/he blocked out the entire event? 11. Has s/he lost interest in doing things that s/he used to like to do since the trauma(s)? 12. Since the trauma(s), does your child show a restricted range of positive 0 1 2 3 4		heart racing, s	shaking hands, sweaty	, short of breath, or si	ck to his/her					
trauma(s)? For example, if other people talk about what happened, does s/he walk away or change the topic? 9. Does your child try to avoid things or places that remind him/her of the trauma(s)? For example, a child who was in a car wreck might try to avoid getting into a car. Or, a child who was in a flood might tell you not to drive over a bridge. Or, a child who saw domestic violence might be nervous to go in the house where it occurred. Or, a girl who was sexually abused might be nervous about going to bed because that's where she was abused before. 10. Does your child have difficulty remembering the whole incident? Has s/he blocked out the entire event? 11. Has s/he lost interest in doing things that s/he used to like to do since the trauma(s)? 12. Since the trauma(s), does your child show a restricted range of positive 0 1 2 3 4										
walk away or change the topic? 9. Does your child try to avoid things or places that remind him/her of the trauma(s)? For example, a child who was in a car wreck might try to avoid getting into a car. Or, a child who was in a flood might tell you not to drive over a bridge. Or, a child who saw domestic violence might be nervous to go in the house where it occurred. Or, a girl who was sexually abused might be nervous about going to bed because that's where she was abused before. 10. Does your child have difficulty remembering the whole incident? Has s/he blocked out the entire event? 11. Has s/he lost interest in doing things that s/he used to like to do since the trauma(s)? 12. Since the trauma(s), does your child show a restricted range of positive 0 1 2 3 4	8.	Does your chi	ild try to avoid conver	sations that might ren	nind him/her of the	0	1	2	3	4
9. Does your child try to avoid things or places that remind him/her of the trauma(s)? For example, a child who was in a car wreck might try to avoid getting into a car. Or, a child who was in a flood might tell you not to drive over a bridge. Or, a child who saw domestic violence might be nervous to go in the house where it occurred. Or, a girl who was sexually abused might be nervous about going to bed because that's where she was abused before. 10. Does your child have difficulty remembering the whole incident? Has s/he blocked out the entire event? 11. Has s/he lost interest in doing things that s/he used to like to do since the trauma(s)? 12. Since the trauma(s), does your child show a restricted range of positive 0 1 2 3 4		trauma(s)? Fo	or example, if other pe	ople talk about what h	nappened, does s/he					
trauma(s)? For example, a child who was in a car wreck might try to avoid getting into a car. Or, a child who was in a flood might tell you not to drive over a bridge. Or, a child who saw domestic violence might be nervous to go in the house where it occurred. Or, a girl who was sexually abused might be nervous about going to bed because that's where she was abused before. 10. Does your child have difficulty remembering the whole incident? Has s/he blocked out the entire event? 11. Has s/he lost interest in doing things that s/he used to like to do since the trauma(s)? 12. Since the trauma(s), does your child show a restricted range of positive 0 1 2 3 4		walk away or	change the topic?							
getting into a car. Or, a child who was in a flood might tell you not to drive over a bridge. Or, a child who saw domestic violence might be nervous to go in the house where it occurred. Or, a girl who was sexually abused might be nervous about going to bed because that's where she was abused before. 10. Does your child have difficulty remembering the whole incident? Has s/he blocked out the entire event? 11. Has s/he lost interest in doing things that s/he used to like to do since the trauma(s)? 12. Since the trauma(s), does your child show a restricted range of positive 0 1 2 3 4	9.	Does your chi	ild try to avoid things	or places that remind	him/her of the	0	1	2	3	4
over a bridge. Or, a child who saw domestic violence might be nervous to go in the house where it occurred. Or, a girl who was sexually abused might be nervous about going to bed because that's where she was abused before. 10. Does your child have difficulty remembering the whole incident? Has s/he blocked out the entire event? 11. Has s/he lost interest in doing things that s/he used to like to do since the trauma(s)? 12. Since the trauma(s), does your child show a restricted range of positive 0 1 2 3 4		trauma(s)? Fo	or example, a child wh	o was in a car wreck	might try to avoid					
go in the house where it occurred. Or, a girl who was sexually abused might be nervous about going to bed because that's where she was abused before. 10. Does your child have difficulty remembering the whole incident? Has s/he blocked out the entire event? 11. Has s/he lost interest in doing things that s/he used to like to do since the trauma(s)? 12. Since the trauma(s), does your child show a restricted range of positive 13. 4		getting into a	car. Or, a child who v	vas in a flood might te	ell you not to drive					
be nervous about going to bed because that's where she was abused before. 10. Does your child have difficulty remembering the whole incident? Has s/he blocked out the entire event? 11. Has s/he lost interest in doing things that s/he used to like to do since the trauma(s)? 12. Since the trauma(s), does your child show a restricted range of positive 13. 4		over a bridge.	Or, a child who saw	domestic violence mig	ght be nervous to					
10. Does your child have difficulty remembering the whole incident? Has s/he blocked out the entire event? 11. Has s/he lost interest in doing things that s/he used to like to do since the trauma(s)? 12. Since the trauma(s), does your child show a restricted range of positive 13. 4		go in the hous	se where it occurred. (Or, a girl who was sex	ually abused might					
blocked out the entire event? 11. Has s/he lost interest in doing things that s/he used to like to do since the trauma(s)? 12. Since the trauma(s), does your child show a restricted range of positive 13. The property of the trauma(s) of the										
11. Has s/he lost interest in doing things that s/he used to like to do since the trauma(s)? 12. Since the trauma(s), does your child show a restricted range of positive 13. 4	10.			nembering the whole i	ncident? Has s/he		_			
trauma(s)? 12. Since the trauma(s), does your child show a restricted range of positive 0 1 2 3 4										
12. Since the trauma(s), does your child show a restricted range of positive 0 1 2 3 4	11.	Has s/he lost	interest in doing thing	s that s/he used to like	e to do since the	0	1	2	3	4
		trauma(s)?								
emotions on his/her face compared to before?	12.	Since the trau	ma(s), does your child	d show a restricted rar	nge of positive	0	1	2	3	4
		emotions on h	nis/her face compared	to before?						

Client Initials:	Client ID:	Date of Completion: /	/

YCPC (Caregiver: English) continued

13.	Has your child lost hope for the future? For example, s/he believes will not					
14.	have fun tomorrow, or will never be good at anything. Since the trauma(s) has your child become more distant and withdrawn	0	1	2	3	4
14.	from family members, relatives, or friends?	U	1	2	3	4
15.	Has s/he had a hard time falling asleep or staying asleep since the	0	1	2	3	4
	trauma(s)?					
16.	Has your child become more irritable, or had outbursts of anger, or	0	1	2	3	4
	developed extreme temper tantrums since the trauma(s)?					
17.	Has your child had more trouble concentrating since the trauma(s)?	0	1	2	3	4
18.	Has s/he been more "on the alert" for bad things to happen? For example,	0	1	2	3	4
	does s/he look around for danger?					
19.	Does your child startle more easily than before the trauma(s)? For example,	0	1	2	3	4
	if there's a loud noise or someone sneaks up behind him/her, does s/he jump					
	or seem startled?					
20.	Has your child become more physically aggressive since the trauma(s)?	0	1	2	3	4
	Like hitting, kicking, biting, or breaking things.					
21.	Has s/he become more clingy to you since the trauma(s)?	0	1	2	3	4
22.	Did night terrors start or get worse after the trauma(s)? Night terrors are	0	1	2	3	4
	different from nightmares: in night terrors a child usually screams in their					
	sleep, they don't wake up, and they don't remember it the next day.					
23.	Since the trauma(s), has your child lost previously acquired skills? For	0	1	2	3	4
	example, lost toilet training? Or, lost language skills? Or, lost motor skills					
	working snaps, buttons, or zippers?					
24.	Since the trauma(s), has your child developed any new fears about things	0	1	2	3	4
	that <u>don't seem related</u> to the trauma(s)? What about going to the bathroom					
	alone? Or, being afraid of the dark?					
	FUNCTIONAL IMPAIRMENT					
	Do the symptoms that you endorsed above get in the way of your child's					
	ability to function in the following areas?					
25.	Do (symptoms) substantially "get in the way" of how s/he gets along with	0	1	2	3	4
	you, interfere in your relationship, or make you feel upset or annoyed?					
26.	Do these (symptoms) "get in the way" of how s/he gets along with brothers	0	1	2	3	4
	or sisters, and make them feel upset or annoyed?					
27.	Do (symptoms) "get in the way" of how s/he gets along with friends at all –	0	1	2	3	4
	at daycare, school, or in your neighborhood?					
28.	Do these (symptoms) "get in the way" with the teacher or the class more	0	1	2	3	4
	than average?					
29.	Do (symptoms) make it harder for you to take him/her out in public than it	0	1	2	3	4
	would be with an average child?					
	Is it harder to go out with your child to places like the grocery store? Or to a					
	restaurant?				_	
30.	Do you think that these behaviors cause your child to feel upset?	0	1	2	3	4

[©] Michael Scheeringa, MD, MPH, 2010, Tulane University, New Orleans, LA. <u>mscheer@tulane.edu</u>. This form may be reproduced and used for free, but not sold, without further permission from the author.

Young Child PTSD Checklist Caregiver Response Scale

Parent Proxy Anxiety - Short Form 8a

Please respond to each question or statement by marking one box per row.

In the past 7 days	Never	Almost Never	Sometimes	Often	Almost Always
My child felt nervous	1	2	3	4	5
My child felt scared	1	2	3	4	5
My child felt worried	1	2	3	4	5
My child felt like something awful might happen	1	2	3	4	5
My child worried when he/she was at home	1	2	3	4	5
My child got scared really easy	1	2	3	4	5
My child worried about what could happen to him/her	1	2	3	4	5
My child worried when he/she went to bed at night	1	2	3	4	5

Response Scale for PROMIS

1 Never

2 Almost 3
Sometimes

4
Often

5
Almost
Always

Client Initials:	Client ID:	Da	Date of Completion://						
SHORT MOOD AND FEELINGS QUESTIONNAIRE (Caregiver: English)									
I'm going to ask you some questions about how your child might have been feeling or acting recently.									
For each question, please answer how much your child has felt or acted this way <u>in the past two</u> <u>weeks</u> .									
If a sentence was true about your child most of the time, check TRUE. If it was only sometimes true, check SOMETIMES. If a sentence was not true about your child, check NOT TRUE									
		True	Sometimes	Not True					
		2	1	0					
1. S/he felt	miserable or unhappy.								
2. S/he did:	n't enjoy anything at all.								
3. S/he felt nothing.	so tired s/he just sat around and did								
4. S/he was	very restless.								
5. S/he felt	s/he was no good any more.								
6. S/he crie	d a lot.								
7. S/he four concentr	nd it hard to think properly or ate.								
8. S/he hate	ed him/herself.								
9. S/he felt	s/he was a bad person.								
10. S/he felt	lonely.								

Copyright Adrian Angold & Elizabeth J. Costello, 1987; Developmental Epidemiology Program; Duke University

11. S/he thought nobody really loved him/her.

12. S/he thought s/he could never be as good as

13. S/he felt s/he did everything wrong.

other kids.

Response Scale for SMFQ

0 1 2
Not True Sometimes True