



EBP INTAKE ASSESSMENT PACKET

MATCH-ADTC

Ages 0-4 Years English

| Required Forms |
|--|
| 1. Demographic Information: |
| Client Intake Face Sheet □ |
| |
| 2. Child's Top Problems: |
| Top Problems Assessment- Caregiver Report □ |
| 3. Child's Trauma History: |
| _ |
| Trauma History Screen- Caregiver Report □ |
| 4. Child's Behavior & Functioning: |
| <i>PPSC</i> - Caregiver Report □ |
| Supplemental Assessments |
| (Included in Packet) |
| Child Depression: SMFQ- Caregiver Report |
| Child Anxiety produce control of the |
| Child Anxiety: <i>PROMIS</i> - Caregiver Report |
| Child Trauma Symptoms: YCPC- Caregiver Report |
| Supplemental Assessments |
| (Included in Supplemental Assessment Packet) |
| Caregiver Symptoms: |
| PSS (Caregiver Stress) |
| PCL-5 (Caregiver Trauma Symptoms) |
| CESD-R (Caregiver Depression) |



Intake Facesheet



VALIDATION REQUIREMENTS AND SYMBOLS EXPLAINED

- ! This symbol means the field is one of the minimum fields that must be filled out to save the record. No data will be saved unless these fields are completed.
- This symbol means the field is a required field in order to save the record as completed. Although you can save the record, the system does not consider the record to be completed unless ALL of these fields are completed.

| Direct Service Provider User Information | | | | | | | | | | | | |
|---|--------|--------------------------------------|-----------|-----------------|-----------|---------------------------------|--------------|---|---|--|--|--|
| Clinician First and Last Name: | | | | | | | | | | | | |
| Treatment Setting: Circle only ONE | | | | Hospital Resid | | | | ych Residential Treatment Facility Shelt sidential Treatment Center Trair hool-Based Othe | | | | |
| | | | Child | ild Information | | | | | | | | |
| First Initial Child's First Name: | | | | First I | Initial (| Child's Last Nam | ie: <u>I</u> | | | | | |
| Date of Birth: ! | | | | Age: | | | | | | | | |
| Sex: ! | П | Female | | _ | Interse | х | | | | | | |
| | | Male | | | Other (| specify)→ | | | | | | |
| Grade (current): * | | | | | | | | | | | | |
| Race: * | _ | American Indian Native | or Alaska | _ | Black o | r African American | | _ | White | | | |
| | | Asian | | 0 | | Hawaiian or Other Islander | | _ | Other (specify) | | | |
| Hispanic Origin: * | | Yes, Cuban | | | Yes, of | Hispanic/Latino Orig | gin | | Yes, South or Central American | | | |
| | | Yes, Mexican, Me American, Chicar | | | Yes, Pu | Yes, Puerto Rican | | | No, Not of Hispanic, Latino, or Spanish Origin | | | |
| City/town: | | | | ST: | | | Z * | ip: | | | | |
| | | Ch | ild Ide | ntifica | ation | Codes | | | | | | |
| Agency-assigned Client ID Number (not PHI): ! | | | | PSDC | RS Clie | nt ID Number: ! | ! | | | | | |
| | | | Family | y Info | rmati | on | | | | | | |
| Caregiver 1 Relationship: * | | | | Careg | iver 2 | Relationship: | | | | | | |
| Preferred Language of Adult Participating in Treatment: * | | | | | | | | | | | | |
| Does the adult participating in t | reatmo | ent speak Engli | ish? | | Yes | | | | No | | | |
| Primary Language of Child: | | | | | | | • | ' | | | | |
| Family Composition: * Select the choice that best describes | | Two parent famil | у | | | parent - cal/adoptive parent | | | Relative/guardian | | | |
| the composition of the family. | | Single Parent wit unrelated partne | | | Blende | d Family | | | Other | | | |



Intake Facesheet



| Living Situation of Child: * | | College Dormitory | | Job Corps | | Psychiatric Hospital |
|---|----------|--|------------------------------|---|---|---|
| What is the child's living situation? | | Crisis Residence | | Medical Hospital | | Residential Treatment Facility |
| | | DCF Foster Home | | Mentor | | TFC Foster Home (privately licensed) |
| | | Group Home | | Military Housing | | Transitional Housing |
| | | Homeless/Shelter | | Other (specify): | | |
| | | Jail/Correctional Facility | | Private Residence | | |
| | | System | Invo | olvement | | |
| Child/Family involved with DCF? | * | | | Yes | | No |
| If child / family is involved with | DCF, p | lease complete ALL of t | he fol | lowing questions: | | |
| DCF Case ID: (if available) | | | _ | Person Link ID: vailable) | | |
| | | Child Protective Services – In-Home | | Family with Service Needs – (FWSN) In-Home | | Not DCF – On Probation |
| DCF Status: | 0 | Child Protective Services – Out of Home | | Family with Service Needs (FWSN) Out of Home | _ | Not DCF – Other Court Involved |
| Dei Status. | 0 | Dual Commitment (JJ and Child Protective Services) | | Juvenile Justice (delinquency) commitment | П | Termination of Parental Rights |
| | | Family Assessment Response | | Not DCF | | Voluntary Services Program |
| DCF Regional Office: | | | | | | |
| Youth involved with Juvenile Jus | stice (J | J) System? * | | Yes | | No |
| If youth is involved with JJ, pleas | se com | plete ALL of the followi | ing qu | estions: | | |
| CSSD Client ID: (if available) | | | CSSD Case ID: (if available) | | | |
| CSSD Case Type: | | | | Delinquency | | Family with Service Needs (Status Offense) |
| | | Administrative Supervision | | Juvenile probation | _ | Restore Probation |
| CSSD Case Status: | 0 | Extended Probation | | Non-Judicial FWSN Family Service Agreement | _ | Suspended Order |
| coop case status. | | Interim Orders | | Non-Judicial Supervision (NJS) | | Waived PDS - Probation |
| | | Judicial FWSN Supervision | | Non-Judicial Supervision Agreement | | |
| Court District: | | | | | | |
| Court Handling Decision: | | | | Judicial | _ | Non-Judicial |
| | | Specific Trea | tmei | nt Information | | |
| What treatment model are you | using v | with this child? * | | TF-CBT | | MATCH-ADTC |
| First Clinical Session Date: * Date of first EBP clinical session | | | | | | |



Intake Facesheet



| Treatment Information | | | | | | | | |
|---|-----------|--|-------|--|--------|--------------------------------|--|--|
| Agency Referral Date/Request for Service: * Date child was referred to agency | | | | is the intake date for the client at tency? | | | | |
| Referral Date: * Date referred for EBP services | | | Inta | ke Date: EBP Intake Date | | | | |
| Referral Source: * Select the source of the EBP referral | | Child Youth-Family Support Center (CYFSC) | | Family Advocate | | Physician | | |
| | | Community Natural Support | | Foster Parent | 0 | Police | | |
| | | Congregate Care Facility | | Info-Line (211) | | Probation/Court | | |
| | | CTBHP/Insurer | | Juvenile Probation / Court | | Psychiatric Hospital | | |
| | | DCF | | Other Community Provider Agency | | School | | |
| | | Detention Involved | | Other Program within Agency | | Self/Family | | |
| | | Emergency Department | | Other State Agency | | | | |
| Assessment Outcome: What was the outcome of the referral to | | Assessment not completed | | Not appropriate for selected EBP | | No treatment needed | | |
| the agency's EBP team? * | | Appropriate for selected EBP | | Not appropriate for selected EBP but needs other treatment | | | | |
| CGI: Considering your experie | nce, h | ow severe are the child' | s emo | tional, behavioral, and/or c | ogniti | ive concerns at the time of | | |
| Intake? Circle only ONE:* | | | | | | Among the most severe symptoms | | |
| Normal Slightly Severe M | lildly S | evere Moderately Seve | ere | Markedly Severe Very Sev | ere | that any child may experience | | |
| | | Treatment I | nfor | mation: School | | | | |
| During the 3 months prior to the start of | f EBP tre | eatment | | | | | | |
| Child's school attendance: * | | Good (few or no days missed) | | No School Attendance: Child Too Young for School | | No School Attendance: Other | | |
| | | Fair (several days missed) | | No School Attendance: Child Suspended/Expelled from School | | | | |
| | | Poor (many days missed) | | No School Attendance: Child Dropped Out of School | | | | |
| Suspended or expelled: * | | | | Yes | | No | | |
| IEP: *Does the child have an Individual | Educati | on Plan (special education)? | | Yes | | No | | |
| | | Treatment | Infor | mation: Legal | | | | |
| During the 3 months prior to the start of | f EBP tre | | | | | | | |
| Arrested: * Has the child been arrest | ed since | start of treatment? | | Yes | | No | | |
| Detained or incarcerated: * Has incarcerated since start of treatment? | the chile | d been detained or | | Yes | | No | | |
| | | Treatment In | form | ation: Medical | | | | |
| During the 3 months prior to the start of | f EBP tre | eatment | | | | | | |
| Alcohol and/or drugs problems: | * | | | Yes | | No | | |
| Evaluated in ER/ED for psychiate | ric issu | es: * | | Yes | | No | | |
| Certified medically complex: * | | | П | Yes | | No | | |

Client ID:

Date of Completion: ___/___/___

Copyright © 2015, President and Fellows of Harvard College. All Rights Reserved.

Client Initials: _____

1

2

3

These copyrighted works were created by John R. Weisz, Ph.D., Kristel Thomassin, Ph.D., Jacqueline Hersh, Ph.D., and Rachel Vaughn-Coaxum, M.A., of Harvard's Laboratory for Youth Mental Health.

| Client Initials: | Client ID: | Date of Completion: | / / | / |
|------------------|------------|---------------------|-----|---|
| | | | , | |



PPSC (Caregiver: English)

18 months, 0 days to 65 months, 31 days *V1.06, 9-1-16*

| PRESCHOOL PE | EDIATRIC SYMPTOM (| CHECKLIST | (PPSC |
|--------------|--------------------|-----------|-------|
|--------------|--------------------|-----------|-------|

These questions are about your child's behavior. Think about what you would expect of other children the same age, and tell us how much each statement applies to your child.

| | | Not at all | Somewnat | very wuch |
|-----------------|---|------------|----------|-----------|
| Does your child | Seem nervous or afraid? · · · · · · · · · | 0 | 1 | 2 |
| | Seem sad or unhappy? · · · · · · · · | . (0) | 1 | 2 |
| | Get upset if things are not done in a certainway? · | . (0) | 1 | 2 |
| | Have a hard time with change? · · · · · · | . (0) | 1 | 2 |
| | Have trouble playing with other children? · · · | . ① | 1 | 2 |
| | Break things on purpose? · · · · · · · | . ① | 1 | 2 |
| | Fight with other children? · · · · · · · | • | 1 | 2 |
| | Have trouble paying attention? · · · · · · | . (0) | 1 | 2 |
| | Have a hard time calming down? · · · · · | • | 1 | 2 |
| | Have trouble staying with one activity? · · · · | . (0) | 1 | 2 |
| ls your child | Aggressive? · · · · · · · · · · · | . (0) | 1 | 2 |
| | Fidgety or unable to sit still? · · · · · · · | . (0) | 1 | 2 |
| | Angry? · · · · · · · · · · · | . (0) | 1 | 2 |
| Is it hard to | Take your child out in public? · · · · · · | • | 1 | 2 |
| | Comfort your child? · · · · · · · · · | . (0) | 1 | 2 |
| | Know what your child needs? · · · · · · | . ① | 1 | 2 |
| | Keep your child on a schedule or routine? · · · | . (0) | 1 | 2 |
| | Get your child to obey you? · · · · · · · | • | 1 | 2 |
| | | | | |



copyright © Tufts Medical Center, 2010. All rights reserved.

Response Scale for PPSC

0 1 2 Not at all Somewhat Very Much

| | | | _ | _ |
|-------------------|------------|---------------------|-----|---|
| Client Initials: | Client ID: | Date of Completion: | / | / |
| CIICITE IIIICIAI3 | CIICITE ID | Date of Completion. | _// | / |

Trauma History Screen (THS) (Caregiver: English)

| | Directions: Ask how many times each event happened, and how much it affected the child when it happened and now. | How many times has this happened? | | | The worst time this happened, how much did it affect him/her? | | | | | How much does this still affect your child? | | | | | | |
|--------|--|-----------------------------------|------|-----------|---|-----------|------------|--------------|------------|---|-----------|------------|--------------|------------|-------------|-----------|
| | "Has your child ever" | Never | Once | 2-3 times | 4-10 times | 10+ times | Not at all | A little bit | Moderately | Quite a bit | Extremely | Not at all | A little bit | Moderately | Quite a bit | Extremely |
| 1 | Been in or seen a very bad accident? | | | | | | 1 | 2 | 3 | 4 | 5 | 1 | 2 | 3 | 4 | 5 |
| 2 | Had someone s/he know been so badly injured or sick that s/he almost died? | | | | | | 1 | 2 | 3 | 4 | 5 | 1 | 2 | 3 | 4 | 5 |
| 3 | Known somebody who died? | | | | | | 1 | 2 | 3 | 4 | 5 | 1 | 2 | 3 | 4 | 5 |
| 4 | Been so sick or hurt that you or the doctor thought s/he might die? | | | | | | 1 | 2 | 3 | 4 | 5 | 1 | 2 | 3 | 4 | 5 |
| 5 | Been unexpectedly separated from someone who s/he depends on for love or security for more than a few days? | | | | | | 1 | 2 | 3 | 4 | 5 | 1 | 2 | 3 | 4 | 5 |
| 6 | Had somebody close to him/her try to kill or hurt themself? | | | | | | 1 | 2 | 3 | 4 | 5 | 1 | 2 | 3 | 4 | 5 |
| 7 | Been physically hurt or threatened by someone? | | | | | | 1 | 2 | 3 | 4 | 5 | 1 | 2 | 3 | 4 | 5 |
| 8 | Been robbed or seen someone get robbed? | | | | | | 1 | 2 | 3 | 4 | 5 | 1 | 2 | 3 | 4 | 5 |
| 9 | Been kidnapped by somebody? | | | | | | 1 | 2 | 3 | 4 | 5 | 1 | 2 | 3 | 4 | 5 |
| 1 0 | Been in or seen a hurricane, earthquake, tornado, or bad fire? | | | | | | 1 | 2 | 3 | 4 | 5 | 1 | 2 | 3 | 4 | 5 |
| 1 | Been attacked by a dog or other animal? | | | | | | 1 | 2 | 3 | 4 | 5 | 1 | 2 | 3 | 4 | 5 |
| 1 2 | Seen or heard people physically fighting or threatening to hurt each other? | | | | | | 1 | 2 | 3 | 4 | 5 | 1 | 2 | 3 | 4 | 5 |
| 1 3 | Seen or heard somebody shooting a gun, using a knife, or using another weapon? | | | | | | 1 | 2 | 3 | 4 | 5 | 1 | 2 | 3 | 4 | 5 |
| 1 4 | Seen a family member arrested or in jail? | | | | | | 1 | 2 | 3 | 4 | 5 | 1 | 2 | 3 | 4 | 5 |
| 1 5 | Had a time in your life when s/he did not have the right care (e.g. food, clothing, a place to live)? | | | | | | 1 | 2 | 3 | 4 | 5 | 1 | 2 | 3 | 4 | 5 |
| 1 6 | Been forced to see or do something sexual? | | | | | | 1 | 2 | 3 | 4 | 5 | 1 | 2 | 3 | 4 | 5 |
| 1 7 | Seen or heard someone else being forced to do something sexual? | | | | | | 1 | 2 | 3 | 4 | 5 | 1 | 2 | 3 | 4 | 5 |
| 1 8 | Watched people using drugs (like smoking, sniffing, or using needles)? | | | | | | 1 | 2 | 3 | 4 | 5 | 1 | 2 | 3 | 4 | 5 |
| 1 9 | Seen something else that was very scary or where s/he thought somebody might get hurt or die? | | | | | | 1 | 2 | 3 | 4 | 5 | 1 | 2 | 3 | 4 | 5 |

20. Which one **bothers your child the MOST** right now: #_____ How long ago did it happen: _____

Response Scale for THS

1 2 3 4 5

Not at All Bit Moderately Quite Extremely A bit

| Client Initials: | Client ID: | Date of Completion: / | / |
|------------------|------------|-----------------------|---|

YOUNG CHILD PTSD CHECKLIST (YCPC) (Caregiver: English)

For Child under 7 years old.

Below is a list of symptoms that children can have after life-threatening events.

When you think of ALL the life-threatening traumatic events from the first page, circle the number below (0-4) that best describes how often the symptom has bothered you in the LAST 2 WEEKS.

| Not at all Once in a while 2 to 4 times a week/ Malmost always Everyday | | 0 1 2 3 | | | | | 4 | | | | | |
|--|-----|-----------------|--------------------------|--------------------------|----------------------|------|---|------|------|-------------|--|--|
| 1. Does your child have intrusive memories of the trauma? Does s/he bring it up on his/her own? 0 1 2 3 4 2. Does your child re-enact the trauma in play with dolls or toys? This would be seenes that look just like the trauma. Or does s/he act it out by him/herself or with other kids? 0 1 2 3 4 3. Is your child having more nightmares since the trauma(s) occurred? 0 1 2 3 4 4. Does your child act like the traumatic event is happening to him/her again, even when it isn't? This is where a child is acting like they are back in the traumatic event and aren't in touch with reality. This is a pretty obvious thing when it happens. 0 1 2 3 4 5. Since the trauma(s) has s/he had episodes when s/he seems to freeze? You may have tried to snap him/her out of it but s/he was unresponsive. 0 1 2 3 4 6. Does s/he get upset when exposed to reminders of the event(s)? For example, a child who was in a car wreck might be nervous when it is raining. Or, a child who saw domestic violence might be nervous when it is raining. Or, a child who saw domestic violence might be nervous when other people argue. Or, a girl who was sexually abused might be nervous when other people alk about what happened, does s/he walk away or change the topic? 0 1 2 3 4 8. Does your c | | Not at all | | 2 to 4 times a week/ | | eek/ | | Ever | yday | | | |
| Up on his/her own? 2. Does your child re-enact the trauma in play with dolls or toys? This would be scenes that look just like the trauma. Or does s/he act it out by him/herself or with other kids? 3. Is your child having more nightmares since the trauma(s) occurred? 0 | | | Once in a while | Half the time | Almost always | | | | | | | |
| Up on his/her own? 2. Does your child re-enact the trauma in play with dolls or toys? This would be scenes that look just like the trauma. Or does s/he act it out by him/herself or with other kids? 3. Is your child having more nightmares since the trauma(s) occurred? 0 | | T = | | | | | | 1 _ | 1 - | | | |
| Does your child re-enact the trauma in play with dolls or toys? This would be scenes that look just like the trauma. Or does s/he act it out by him/herself or with other kids? 3. Is your child having more nightmares since the trauma(s) occurred? 0 1 2 3 4 | 1. | - | | Does s/he bring it | 0 | 1 | 2 | 3 | 4 | | | |
| be scenes that look just like the trauma. Or does s/he act it out by him/herself or with other kids? 3. Is your child having more nightmares since the trauma(s) occurred? 4. Does your child act like the traumatic event is happening to him/her again, even when it isn't? This is where a child is acting like they are back in the traumatic event and aren't in touch with reality. This is a pretty obvious thing when it happens. 5. Since the trauma(s) has s/he had episodes when s/he seems to freeze? You may have tried to snap him/her out of it but s/he was unresponsive. 6. Does s/he get upset when exposed to reminders of the event(s)? For example, a child who was in a car wreck might be nervous when it is raining. Or, a child who was in a hurricane might be nervous when other people argue. Or, a girl who was sexually abused might be nervous when someone touches her. 7. Does your child get physically distressed when exposed to reminders? Like heart racing, shaking hands, sweaty, short of breath, or sick to his/her stomach? Think of the same type of examples as in #6. 8. Does your child try to avoid conversations that might remind him/her of the trauma(s)? For example, if other people talk about what happened, does s/he walk away or change the topic? 9. Does your child try to avoid things or places that remind him/her of the trauma(s)? For example, a child who was in a car wreck might try to avoid getting into a car. Or, a child who was in a flood might tell you not to drive over a bridge. Or, a child who was in a flood might tell you not to drive over a bridge. Or, a child who was in a flood might tell you not to drive over a bridge. Or, a child who saw domestic violence might be nervous to go in the house where it occurred. Or, a girl who was sexually abused might be nervous about going to bed because that's where she was abused before. 10. Does your child have difficulty remembering the whole incident? Has s/he blocked out the entire event? 11. Has s/he lost interest in doing things that s/he used to like to | | 1 1 | | | 0.001 | - | | | | | | |
| him/herself or with other kids? Is your child having more nightmares since the trauma(s) occurred? 0 | 2. | _ | | | = | 0 | 1 | 2 | 3 | 4 | | |
| 3. Is your child having more nightmares since the trauma(s) occurred? 4. Does your child act like the traumatic event is happening to him/her again, even when it isn't? This is where a child is acting like they are back in the traumatic event and aren't in touch with reality. This is a pretty obvious thing when it happens. 5. Since the trauma(s) has s/he had episodes when s/he seems to freeze? You may have tried to snap him/her out of it but s/he was unresponsive. 6. Does s/he get upset when exposed to reminders of the event(s)? For example, a child who was in a car wreck might be nervous while riding in a car now. Or, a child who saw domestic violence might be nervous when other people argue. Or, a girl who was sexually abused might be nervous when other people argue. Or, a girl who was sexually abused might be nervous when other people argue. Or, a child who saw domestic violence might be nervous when other people argue. Or, a girl who was sexually abused might be nervous when other people argue. Or, a child who saw domestic violence might be nervous when other people argue. Or, a child voice of breath, or sick to his/her stomach? Think of the same type of examples as in #6. 8. Does your child try to avoid conversations that might remind him/her of the trauma(s)? For example, if other people talk about what happened, does s/he walk away or change the topic? 9. Does your child try to avoid things or places that remind him/her of the trauma(s)? For example, a child who was in a car wreck might try to avoid getting into a car. Or, a child who was in a flood might tell you not to drive over a bridge. Or, a child who saw domestic violence might be nervous to go in the house where it occurred. Or, a girl who was sexually abused might be nervous about going to bed because that's where she was abused before. 10. Does your child have difficulty remembering the whole incident? Has s/he blocked out the entire event? 11. Has s/he lost interest in doing thing | | | | ıma. Or does s/he act | it out by | | | | | | | |
| 4. Does your child act like the traumatic event is happening to him/her again, even when it isn't? This is where a child is acting like they are back in the traumatic event and aren't in touch with reality. This is a pretty obvious thing when it happens. 5. Since the trauma(s) has s/he had episodes when s/he seems to freeze? You may have tried to snap him/her out of it but s/he was unresponsive. 6. Does s/he get upset when exposed to reminders of the event(s)? For example, a child who was in a car wreck might be nervous while riding in a car now. Or, a child who was in a hurricane might be nervous when other people argue. Or, a girl who was sexually abused might be nervous when someone touches her. 7. Does your child get physically distressed when exposed to reminders? Like heart racing, shaking hands, sweaty, short of breath, or sick to his/her stomach? Think of the same type of examples as in #6. 8. Does your child try to avoid conversations that might remind him/her of the trauma(s)? For example, if other people talk about what happened, does s/he walk away or change the topic? 9. Does your child try to avoid things or places that remind him/her of the trauma(s)? For example, a child who was in a flood might tell you not to drive over a bridge. Or, a child who saw domestic violence might be nervous to go in the house where it occurred. Or, a girl who was sexually abused might be nervous about going to bed because that's where she was abused before. 10. Does your child have difficulty remembering the whole incident? Has s/he blocked out the entire event? 11. Has s/he lost interest in doing things that s/he used to like to do since the trauma(s)? 12. Since the trauma(s), does your child show a restricted range of positive 0 1 2 3 4 | | | | | | | | | | | | |
| even when it isn't? This is where a child is acting like they are back in the traumatic event and aren't in touch with reality. This is a pretty obvious thing when it happens. 5. Since the trauma(s) has s/he had episodes when s/he seems to freeze? You may have tried to snap him/her out of it but s/he was unresponsive. 6. Does s/he get upset when exposed to reminders of the event(s)? For example, a child who was in a car wreck might be nervous while riding in a car now. Or, a child who saw domestic violence might be nervous when other people argue. Or, a girl who was sexually abused might be nervous when someone touches her. 7. Does your child get physically distressed when exposed to reminders? Like heart racing, shaking hands, sweaty, short of breath, or sick to his/her stomach? Think of the same type of examples as in #6. 8. Does your child try to avoid conversations that might remind him/her of the trauma(s)? For example, if other people talk about what happened, does s/he walk away or change the topic? 9. Does your child try to avoid things or places that remind him/her of the trauma(s)? For example, a child who was in a car wreck might try to avoid getting into a car. Or, a child who was in a flood might tell you not to drive over a bridge. Or, a child who saw domestic violence might be nervous to go in the house where it occurred. Or, a girl who was sexually abused might be nervous about going to bed because that's where she was abused before. 10. Does your child have difficulty remembering the whole incident? Has s/he blocked out the entire event? 11. Has s/he lost interest in doing things that s/he used to like to do since the trauma(s)? 12. Since the trauma(s), does your child show a restricted range of positive 0 1 2 3 4 | | | | | | | 1 | | | | | |
| traumatic event and aren't in touch with reality. This is a pretty obvious thing when it happens. 5. Since the trauma(s) has s/he had episodes when s/he seems to freeze? You may have tried to snap him/her out of it but s/he was unresponsive. 6. Does s/he get upset when exposed to reminders of the event(s)? For example, a child who was in a car wreck might be nervous while riding in a car now. Or, a child who saw domestic violence might be nervous when other people argue. Or, a girl who was sexually abused might be nervous when someone touches her. 7. Does your child get physically distressed when exposed to reminders? Like heart racing, shaking hands, sweaty, short of breath, or sick to his/her stomach? Think of the same type of examples as in #6. 8. Does your child try to avoid conversations that might remind him/her of the trauma(s)? For example, if other people talk about what happened, does s/he walk away or change the topic? 9. Does your child try to avoid things or places that remind him/her of the trauma(s)? For example, a child who was in a car wreck might try to avoid getting into a car. Or, a child who was in a flood might tell you not to drive over a bridge. Or, a child who saw domestic violence might be nervous to go in the house where it occurred. Or, a girl who was sexually abused might be nervous about going to bed because that's where she was abused before. 10. Does your child have difficulty remembering the whole incident? Has s/he blocked out the entire event? 11. Has s/he lost interest in doing things that s/he used to like to do since the trauma(s)? 12. Since the trauma(s), does your child show a restricted range of positive 0 1 2 3 4 | 4. | | | | | 0 | 1 | 2 | 3 | 4 | | |
| thing when it happens. 5. Since the trauma(s) has s/he had episodes when s/he seems to freeze? You may have tried to snap him/her out of it but s/he was unresponsive. 6. Does s/he get upset when exposed to reminders of the event(s)? For example, a child who was in a car wreck might be nervous while riding in a car now. Or, a child who was in a hurricane might be nervous when it is raining. Or, a child who saw domestic violence might be nervous when other people argue. Or, a girl who was sexually abused might be nervous when someone touches her. 7. Does your child get physically distressed when exposed to reminders? Like heart racing, shaking hands, sweaty, short of breath, or sick to his/her stomach? Think of the same type of examples as in #6. 8. Does your child try to avoid conversations that might remind him/her of the trauma(s)? For example, if other people talk about what happened, does s/he walk away or change the topic? 9. Does your child try to avoid things or places that remind him/her of the walk away or change the topic? 9. Does your child try to avoid things or places that remind him/her of the trauma(s)? For example, a child who was in a car wreck might try to avoid getting into a car. Or, a child who was in a flood might tell you not to drive over a bridge. Or, a child who was domestic violence might be nervous to go in the house where it occurred. Or, a girl who was sexually abused might be nervous about going to bed because that's where she was abused before. 10. Does your child have difficulty remembering the whole incident? Has s/he blocked out the entire event? 11. Has s/he lost interest in doing things that s/he used to like to do since the trauma(s)? 12. Since the trauma(s), does your child show a restricted range of positive 0 1 2 3 4 | | | | | - | | | | | | | |
| 5. Since the trauma(s) has s/he had episodes when s/he seems to freeze? You may have tried to snap him/her out of it but s/he was unresponsive. 6. Does s/he get upset when exposed to reminders of the event(s)? For example, a child who was in a car wreck might be nervous while riding in a car now. Or, a child who was in a hurricane might be nervous when it is raining. Or, a child who saw domestic violence might be nervous when other people argue. Or, a girl who was sexually abused might be nervous when someone touches her. 7. Does your child get physically distressed when exposed to reminders? Like heart racing, shaking hands, sweaty, short of breath, or sick to his/her stomach? Think of the same type of examples as in #6. 8. Does your child try to avoid conversations that might remind him/her of the trauma(s)? For example, if other people talk about what happened, does s/he walk away or change the topic? 9. Does your child try to avoid things or places that remind him/her of the value and thing into a car. Or, a child who was in a car wreck might try to avoid getting into a car. Or, a child who was in a flood might tell you not to drive over a bridge. Or, a child who saw domestic violence might be nervous to go in the house where it occurred. Or, a girl who was sexually abused might be nervous about going to bed because that's where she was abused before. 10. Does your child have difficulty remembering the whole incident? Has s/he blocked out the entire event? 11. Has s/he lost interest in doing things that s/he used to like to do since the trauma(s)? 12. Since the trauma(s), does your child show a restricted range of positive 0 1 2 3 4 | | | | with reality. This is a | pretty obvious | | | | | | | |
| may have tried to snap him/her out of it but s/he was unresponsive. 6. Does s/he get upset when exposed to reminders of the event(s)? For example, a child who was in a car wreck might be nervous while riding in a car now. Or, a child who was in a hurricane might be nervous when it is raining. Or, a child who saw domestic violence might be nervous when other people argue. Or, a girl who was sexually abused might be nervous when someone touches her. 7. Does your child get physically distressed when exposed to reminders? Like heart racing, shaking hands, sweaty, short of breath, or sick to his/her stomach? Think of the same type of examples as in #6. 8. Does your child try to avoid conversations that might remind him/her of the trauma(s)? For example, if other people talk about what happened, does s/he walk away or change the topic? 9. Does your child try to avoid things or places that remind him/her of the trauma(s)? For example, a child who was in a car wreck might try to avoid getting into a car. Or, a child who was in a flood might tell you not to drive over a bridge. Or, a child who saw domestic violence might be nervous to go in the house where it occurred. Or, a girl who was sexually abused might be nervous about going to bed because that's where she was abused before. 10. Does your child have difficulty remembering the whole incident? Has s/he blocked out the entire event? 11. Has s/he lost interest in doing things that s/he used to like to do since the trauma(s)? 12. Since the trauma(s), does your child show a restricted range of positive 0 1 2 3 4 | | | | | | | | | | | | |
| 6. Does s/he get upset when exposed to reminders of the event(s)? For example, a child who was in a car wreck might be nervous while riding in a car now. Or, a child who saw domestic violence might be nervous when other people argue. Or, a girl who was sexually abused might be nervous when someone touches her. 7. Does your child get physically distressed when exposed to reminders? Like heart racing, shaking hands, sweaty, short of breath, or sick to his/her stomach? Think of the same type of examples as in #6. 8. Does your child try to avoid conversations that might remind him/her of the trauma(s)? For example, if other people talk about what happened, does s/he walk away or change the topic? 9. Does your child try to avoid things or places that remind him/her of the trauma(s)? For example, a child who was in a car wreck might try to avoid getting into a car. Or, a child who was in a flood might tell you not to drive over a bridge. Or, a child who saw domestic violence might be nervous to go in the house where it occurred. Or, a girl who was sexually abused might be nervous about going to bed because that's where she was abused before. 10. Does your child have difficulty remembering the whole incident? Has s/he blocked out the entire event? 11. Has s/he lost interest in doing things that s/he used to like to do since the trauma(s)? 12. Since the trauma(s), does your child show a restricted range of positive 0 1 2 3 4 | 5. | | | | | 0 | 1 | 2 | 3 | 4 | | |
| example, a child who was in a car wreck might be nervous while riding in a car now. Or, a child who was in a hurricane might be nervous when it is raining. Or, a child who saw domestic violence might be nervous when other people argue. Or, a girl who was sexually abused might be nervous when someone touches her. 7. Does your child get physically distressed when exposed to reminders? Like heart racing, shaking hands, sweaty, short of breath, or sick to his/her stomach? Think of the same type of examples as in #6. 8. Does your child try to avoid conversations that might remind him/her of the trauma(s)? For example, if other people talk about what happened, does s/he walk away or change the topic? 9. Does your child try to avoid things or places that remind him/her of the trauma(s)? For example, a child who was in a car wreck might try to avoid getting into a car. Or, a child who was in a flood might tell you not to drive over a bridge. Or, a child who saw domestic violence might be nervous to go in the house where it occurred. Or, a girl who was sexually abused might be nervous about going to bed because that's where she was abused before. 10. Does your child have difficulty remembering the whole incident? Has s/he blocked out the entire event? 11. Has s/he lost interest in doing things that s/he used to like to do since the trauma(s)? | | | | | | | | | | | | |
| car now. Or, a child who was in a hurricane might be nervous when it is raining. Or, a child who saw domestic violence might be nervous when other people argue. Or, a girl who was sexually abused might be nervous when someone touches her. 7. Does your child get physically distressed when exposed to reminders? Like heart racing, shaking hands, sweaty, short of breath, or sick to his/her stomach? Think of the same type of examples as in #6. 8. Does your child try to avoid conversations that might remind him/her of the trauma(s)? For example, if other people talk about what happened, does s/he walk away or change the topic? 9. Does your child try to avoid things or places that remind him/her of the trauma(s)? For example, a child who was in a car wreck might try to avoid getting into a car. Or, a child who was in a flood might tell you not to drive over a bridge. Or, a child who saw domestic violence might be nervous to go in the house where it occurred. Or, a girl who was sexually abused might be nervous about going to bed because that's where she was abused before. 10. Does your child have difficulty remembering the whole incident? Has s/he blocked out the entire event? 11. Has s/he lost interest in doing things that s/he used to like to do since the trauma(s)? 12. Since the trauma(s), does your child show a restricted range of positive 0 1 2 3 4 | 6. | | | | | 0 | 1 | 2 | 3 | 4 | | |
| raining. Or, a child who saw domestic violence might be nervous when other people argue. Or, a girl who was sexually abused might be nervous when someone touches her. 7. Does your child get physically distressed when exposed to reminders? Like heart racing, shaking hands, sweaty, short of breath, or sick to his/her stomach? Think of the same type of examples as in #6. 8. Does your child try to avoid conversations that might remind him/her of the trauma(s)? For example, if other people talk about what happened, does s/he walk away or change the topic? 9. Does your child try to avoid things or places that remind him/her of the trauma(s)? For example, a child who was in a car wreck might try to avoid getting into a car. Or, a child who was in a flood might tell you not to drive over a bridge. Or, a child who saw domestic violence might be nervous to go in the house where it occurred. Or, a girl who was sexually abused might be nervous about going to bed because that's where she was abused before. 10. Does your child have difficulty remembering the whole incident? Has s/he blocked out the entire event? 11. Has s/he lost interest in doing things that s/he used to like to do since the trauma(s)? 12. Since the trauma(s), does your child show a restricted range of positive 0 1 2 3 4 | | example, a ch | ild who was in a car v | wreck might be nervou | is while riding in a | | | | | | | |
| other people argue. Or, a girl who was sexually abused might be nervous when someone touches her. 7. Does your child get physically distressed when exposed to reminders? Like heart racing, shaking hands, sweaty, short of breath, or sick to his/her stomach? Think of the same type of examples as in #6. 8. Does your child try to avoid conversations that might remind him/her of the trauma(s)? For example, if other people talk about what happened, does s/he walk away or change the topic? 9. Does your child try to avoid things or places that remind him/her of the trauma(s)? For example, a child who was in a car wreck might try to avoid getting into a car. Or, a child who was in a flood might tell you not to drive over a bridge. Or, a child who saw domestic violence might be nervous to go in the house where it occurred. Or, a girl who was sexually abused might be nervous about going to bed because that's where she was abused before. 10. Does your child have difficulty remembering the whole incident? Has s/he blocked out the entire event? 11. Has s/he lost interest in doing things that s/he used to like to do since the trauma(s)? 12. Since the trauma(s), does your child show a restricted range of positive 0 1 2 3 4 | | | | • | | | | | | | | |
| when someone touches her. 7. Does your child get physically distressed when exposed to reminders? Like heart racing, shaking hands, sweaty, short of breath, or sick to his/her stomach? Think of the same type of examples as in #6. 8. Does your child try to avoid conversations that might remind him/her of the trauma(s)? For example, if other people talk about what happened, does s/he walk away or change the topic? 9. Does your child try to avoid things or places that remind him/her of the trauma(s)? For example, a child who was in a car wreck might try to avoid getting into a car. Or, a child who was in a flood might tell you not to drive over a bridge. Or, a child who saw domestic violence might be nervous to go in the house where it occurred. Or, a girl who was sexually abused might be nervous about going to bed because that's where she was abused before. 10. Does your child have difficulty remembering the whole incident? Has s/he blocked out the entire event? 11. Has s/he lost interest in doing things that s/he used to like to do since the trauma(s)? 12. Since the trauma(s), does your child show a restricted range of positive 0 1 2 3 4 | | raining. Or, a | child who saw domes | tic violence might be | nervous when | | | | | | | |
| 7. Does your child get physically distressed when exposed to reminders? Like heart racing, shaking hands, sweaty, short of breath, or sick to his/her stomach? Think of the same type of examples as in #6. 8. Does your child try to avoid conversations that might remind him/her of the trauma(s)? For example, if other people talk about what happened, does s/he walk away or change the topic? 9. Does your child try to avoid things or places that remind him/her of the trauma(s)? For example, a child who was in a car wreck might try to avoid getting into a car. Or, a child who was in a flood might tell you not to drive over a bridge. Or, a child who saw domestic violence might be nervous to go in the house where it occurred. Or, a girl who was sexually abused might be nervous about going to bed because that's where she was abused before. 10. Does your child have difficulty remembering the whole incident? Has s/he blocked out the entire event? 11. Has s/he lost interest in doing things that s/he used to like to do since the trauma(s)? 12. Since the trauma(s), does your child show a restricted range of positive 0 1 2 3 4 | | | | vas sexually abused m | night be nervous | | | | | | | |
| heart racing, shaking hands, sweaty, short of breath, or sick to his/her stomach? Think of the same type of examples as in #6. 8. Does your child try to avoid conversations that might remind him/her of the trauma(s)? For example, if other people talk about what happened, does s/he walk away or change the topic? 9. Does your child try to avoid things or places that remind him/her of the trauma(s)? For example, a child who was in a car wreck might try to avoid getting into a car. Or, a child who was in a flood might tell you not to drive over a bridge. Or, a child who saw domestic violence might be nervous to go in the house where it occurred. Or, a girl who was sexually abused might be nervous about going to bed because that's where she was abused before. 10. Does your child have difficulty remembering the whole incident? Has s/he blocked out the entire event? 11. Has s/he lost interest in doing things that s/he used to like to do since the trauma(s)? 12. Since the trauma(s), does your child show a restricted range of positive 0 1 2 3 4 | | when someon | e touches her. | | | | | | | | | |
| stomach? Think of the same type of examples as in #6. 8. Does your child try to avoid conversations that might remind him/her of the trauma(s)? For example, if other people talk about what happened, does s/he walk away or change the topic? 9. Does your child try to avoid things or places that remind him/her of the trauma(s)? For example, a child who was in a car wreck might try to avoid getting into a car. Or, a child who was in a flood might tell you not to drive over a bridge. Or, a child who saw domestic violence might be nervous to go in the house where it occurred. Or, a girl who was sexually abused might be nervous about going to bed because that's where she was abused before. 10. Does your child have difficulty remembering the whole incident? Has s/he blocked out the entire event? 11. Has s/he lost interest in doing things that s/he used to like to do since the trauma(s)? 12. Since the trauma(s), does your child show a restricted range of positive 0 1 2 3 4 | 7. | Does your chi | ild get physically distr | ressed when exposed t | o reminders? Like | 0 | 1 | 2 | 3 | 4 | | |
| 8. Does your child try to avoid conversations that might remind him/her of the trauma(s)? For example, if other people talk about what happened, does s/he walk away or change the topic? 9. Does your child try to avoid things or places that remind him/her of the trauma(s)? For example, a child who was in a car wreck might try to avoid getting into a car. Or, a child who was in a flood might tell you not to drive over a bridge. Or, a child who saw domestic violence might be nervous to go in the house where it occurred. Or, a girl who was sexually abused might be nervous about going to bed because that's where she was abused before. 10. Does your child have difficulty remembering the whole incident? Has s/he blocked out the entire event? 11. Has s/he lost interest in doing things that s/he used to like to do since the trauma(s)? 12. Since the trauma(s), does your child show a restricted range of positive 0 1 2 3 4 | | heart racing, s | shaking hands, sweaty | , short of breath, or si | ck to his/her | | | | | | | |
| trauma(s)? For example, if other people talk about what happened, does s/he walk away or change the topic? 9. Does your child try to avoid things or places that remind him/her of the trauma(s)? For example, a child who was in a car wreck might try to avoid getting into a car. Or, a child who was in a flood might tell you not to drive over a bridge. Or, a child who saw domestic violence might be nervous to go in the house where it occurred. Or, a girl who was sexually abused might be nervous about going to bed because that's where she was abused before. 10. Does your child have difficulty remembering the whole incident? Has s/he blocked out the entire event? 11. Has s/he lost interest in doing things that s/he used to like to do since the trauma(s)? 12. Since the trauma(s), does your child show a restricted range of positive 0 1 2 3 4 | | | | | | | | | | | | |
| walk away or change the topic? 9. Does your child try to avoid things or places that remind him/her of the trauma(s)? For example, a child who was in a car wreck might try to avoid getting into a car. Or, a child who was in a flood might tell you not to drive over a bridge. Or, a child who saw domestic violence might be nervous to go in the house where it occurred. Or, a girl who was sexually abused might be nervous about going to bed because that's where she was abused before. 10. Does your child have difficulty remembering the whole incident? Has s/he blocked out the entire event? 11. Has s/he lost interest in doing things that s/he used to like to do since the trauma(s)? 12. Since the trauma(s), does your child show a restricted range of positive 0 1 2 3 4 | 8. | Does your chi | ild try to avoid conver | sations that might ren | nind him/her of the | 0 | 1 | 2 | 3 | 4 | | |
| 9. Does your child try to avoid things or places that remind him/her of the trauma(s)? For example, a child who was in a car wreck might try to avoid getting into a car. Or, a child who was in a flood might tell you not to drive over a bridge. Or, a child who saw domestic violence might be nervous to go in the house where it occurred. Or, a girl who was sexually abused might be nervous about going to bed because that's where she was abused before. 10. Does your child have difficulty remembering the whole incident? Has s/he blocked out the entire event? 11. Has s/he lost interest in doing things that s/he used to like to do since the trauma(s)? 12. Since the trauma(s), does your child show a restricted range of positive 0 1 2 3 4 | | trauma(s)? Fo | or example, if other pe | ople talk about what h | nappened, does s/he | | | | | | | |
| trauma(s)? For example, a child who was in a car wreck might try to avoid getting into a car. Or, a child who was in a flood might tell you not to drive over a bridge. Or, a child who saw domestic violence might be nervous to go in the house where it occurred. Or, a girl who was sexually abused might be nervous about going to bed because that's where she was abused before. 10. Does your child have difficulty remembering the whole incident? Has s/he blocked out the entire event? 11. Has s/he lost interest in doing things that s/he used to like to do since the trauma(s)? 12. Since the trauma(s), does your child show a restricted range of positive 0 1 2 3 4 | | walk away or | change the topic? | | | | | | | | | |
| getting into a car. Or, a child who was in a flood might tell you not to drive over a bridge. Or, a child who saw domestic violence might be nervous to go in the house where it occurred. Or, a girl who was sexually abused might be nervous about going to bed because that's where she was abused before. 10. Does your child have difficulty remembering the whole incident? Has s/he blocked out the entire event? 11. Has s/he lost interest in doing things that s/he used to like to do since the trauma(s)? 12. Since the trauma(s), does your child show a restricted range of positive 0 1 2 3 4 | 9. | Does your chi | ild try to avoid things | or places that remind | him/her of the | 0 | 1 | 2 | 3 | 4 | | |
| over a bridge. Or, a child who saw domestic violence might be nervous to go in the house where it occurred. Or, a girl who was sexually abused might be nervous about going to bed because that's where she was abused before. 10. Does your child have difficulty remembering the whole incident? Has s/he blocked out the entire event? 11. Has s/he lost interest in doing things that s/he used to like to do since the trauma(s)? 12. Since the trauma(s), does your child show a restricted range of positive 0 1 2 3 4 | | trauma(s)? Fo | or example, a child wh | o was in a car wreck | might try to avoid | | | | | | | |
| go in the house where it occurred. Or, a girl who was sexually abused might be nervous about going to bed because that's where she was abused before. 10. Does your child have difficulty remembering the whole incident? Has s/he blocked out the entire event? 11. Has s/he lost interest in doing things that s/he used to like to do since the trauma(s)? 12. Since the trauma(s), does your child show a restricted range of positive 13. 4 | | getting into a | car. Or, a child who v | vas in a flood might te | ell you not to drive | | | | | | | |
| be nervous about going to bed because that's where she was abused before. 10. Does your child have difficulty remembering the whole incident? Has s/he blocked out the entire event? 11. Has s/he lost interest in doing things that s/he used to like to do since the trauma(s)? 12. Since the trauma(s), does your child show a restricted range of positive 13. 4 | | over a bridge. | Or, a child who saw | domestic violence mig | ght be nervous to | | | | | | | |
| 10. Does your child have difficulty remembering the whole incident? Has s/he blocked out the entire event? 11. Has s/he lost interest in doing things that s/he used to like to do since the trauma(s)? 12. Since the trauma(s), does your child show a restricted range of positive 13. 4 | | go in the hous | se where it occurred. (| Or, a girl who was sex | ually abused might | | | | | | | |
| blocked out the entire event? 11. Has s/he lost interest in doing things that s/he used to like to do since the trauma(s)? 12. Since the trauma(s), does your child show a restricted range of positive 13. The property of the trauma(s) of the | | | | | | | | | | | | |
| 11. Has s/he lost interest in doing things that s/he used to like to do since the trauma(s)? 12. Since the trauma(s), does your child show a restricted range of positive 13. 4 | 10. | | | nembering the whole i | ncident? Has s/he | | | | | | | |
| trauma(s)? 12. Since the trauma(s), does your child show a restricted range of positive 0 1 2 3 4 | | blocked out th | ne entire event? | | | | | | | | | |
| 12. Since the trauma(s), does your child show a restricted range of positive 0 1 2 3 4 | 11. | Has s/he lost | interest in doing thing | s that s/he used to like | e to do since the | 0 | 1 | 2 | 3 | 4 | | |
| | | trauma(s)? | | | | | | | | | | |
| emotions on his/her face compared to before? | 12. | Since the trau | ma(s), does your child | d show a restricted rar | nge of positive | 0 | 1 | 2 | 3 | 4 | | |
| | | emotions on h | nis/her face compared | to before? | | | | | | | | |

| Client Initials: | Client ID: | Date of Completion: / | / |
|------------------|------------|-----------------------|---|

YCPC (Caregiver: English) continued

| 13. | Has your child lost hope for the future? For example, s/he believes will not | | | | | |
|-----|---|---|---|---|---|---|
| 14. | have fun tomorrow, or will never be good at anything. Since the trauma(s) has your child become more distant and withdrawn | 0 | 1 | 2 | 3 | 4 |
| 14. | from family members, relatives, or friends? | U | 1 | 2 | 3 | 4 |
| 15. | Has s/he had a hard time falling asleep or staying asleep since the | 0 | 1 | 2 | 3 | 4 |
| | trauma(s)? | | | | | |
| 16. | Has your child become more irritable, or had outbursts of anger, or | 0 | 1 | 2 | 3 | 4 |
| | developed extreme temper tantrums since the trauma(s)? | | | | | |
| 17. | Has your child had more trouble concentrating since the trauma(s)? | 0 | 1 | 2 | 3 | 4 |
| 18. | Has s/he been more "on the alert" for bad things to happen? For example, | 0 | 1 | 2 | 3 | 4 |
| | does s/he look around for danger? | | | | | |
| 19. | Does your child startle more easily than before the trauma(s)? For example, | 0 | 1 | 2 | 3 | 4 |
| | if there's a loud noise or someone sneaks up behind him/her, does s/he jump | | | | | |
| | or seem startled? | | | | | |
| 20. | Has your child become more physically aggressive since the trauma(s)? | 0 | 1 | 2 | 3 | 4 |
| | Like hitting, kicking, biting, or breaking things. | | | | | |
| 21. | Has s/he become more clingy to you since the trauma(s)? | 0 | 1 | 2 | 3 | 4 |
| 22. | Did night terrors start or get worse after the trauma(s)? Night terrors are | 0 | 1 | 2 | 3 | 4 |
| | different from nightmares: in night terrors a child usually screams in their | | | | | |
| | sleep, they don't wake up, and they don't remember it the next day. | | | | | |
| 23. | Since the trauma(s), has your child lost previously acquired skills? For | 0 | 1 | 2 | 3 | 4 |
| | example, lost toilet training? Or, lost language skills? Or, lost motor skills | | | | | |
| | working snaps, buttons, or zippers? | | | | | |
| 24. | Since the trauma(s), has your child developed any new fears about things | 0 | 1 | 2 | 3 | 4 |
| | that <u>don't seem related</u> to the trauma(s)? What about going to the bathroom | | | | | |
| | alone? Or, being afraid of the dark? | | | | | |
| | FUNCTIONAL IMPAIRMENT | | | | | |
| | Do the symptoms that you endorsed above get in the way of your child's | | | | | |
| | ability to function in the following areas? | | | | | |
| 25. | Do (symptoms) substantially "get in the way" of how s/he gets along with | 0 | 1 | 2 | 3 | 4 |
| | you, interfere in your relationship, or make you feel upset or annoyed? | | | | | |
| 26. | Do these (symptoms) "get in the way" of how s/he gets along with brothers | 0 | 1 | 2 | 3 | 4 |
| | or sisters, and make them feel upset or annoyed? | | | | | |
| 27. | Do (symptoms) "get in the way" of how s/he gets along with friends at all – | 0 | 1 | 2 | 3 | 4 |
| | at daycare, school, or in your neighborhood? | | | | | |
| 28. | Do these (symptoms) "get in the way" with the teacher or the class more | 0 | 1 | 2 | 3 | 4 |
| | than average? | | | | | |
| 29. | Do (symptoms) make it harder for you to take him/her out in public than it | 0 | 1 | 2 | 3 | 4 |
| | would be with an average child? | | | | | |
| | Is it harder to go out with your child to places like the grocery store? Or to a | | | | | |
| | restaurant? | | | | _ | |
| 30. | Do you think that these behaviors cause your child to feel upset? | 0 | 1 | 2 | 3 | 4 |

[©] Michael Scheeringa, MD, MPH, 2010, Tulane University, New Orleans, LA. <u>mscheer@tulane.edu</u>. This form may be reproduced and used for free, but not sold, without further permission from the author.

Young Child PTSD Checklist Caregiver Response Scale

Parent Proxy Anxiety - Short Form 8a

Please respond to each question or statement by marking one box per row.

| In the past 7 days | Never | Almost Never | Sometimes | Often | Almost Always |
|---|-------|-----------------|-----------|-------|------------------|
| My child felt nervous | 1 | 2 | 3 | 4 | 5 |
| My child felt scared | 1 | 2 | 3 | 4 | 5 |
| My child felt worried | 1 | 2 | 3 | 4 | 5 |
| My child felt like something awful might happen | 1 | 2 | 3 | 4 | 5 |
| My child worried when he/she was at home | 1 | 2 | 3 | 4 | 5 |
| My child got scared really easy | 1 | 2 | 3 | 4 | 5 |
| My child worried about what could happen to him/her | 1 | 2 | 3 | 4 | 5 |
| My child worried when he/she went to bed at night | 1 | 2 | 3 | 4 | 5 |

Response Scale for PROMIS

1 Never

2 Almost 3
Sometimes

4
Often

5
Almost
Always

| Client Initials: | Client ID: | Da | Date of Completion:// | | | | |
|---|--------------------------------|--------------------|-------------------------|--------------|--|--|--|
| SHORT MOOD AND FEELINGS QUESTIONNAIRE (Caregiver: English) | | | | | | | |
| I'm going to ask you some questions about how your child might have been feeling or acting recently. | | | | | | | |
| For each question, pweeks. | please answer how much your cl | nild has felt or a | cted this way <u>in</u> | the past two | | | |
| If a sentence was true about your child most of the time, check TRUE. If it was only sometimes true, check SOMETIMES. If a sentence was not true about your child, check NOT TRUE | | | | | | | |
| | | True | Sometimes | Not True | | | |
| | | 2 | 1 | 0 | | | |
| 1. S/he felt misera | ble or unhappy. | | | | | | |
| 2. S/he didn't enjo | y anything at all. | | | | | | |
| 3. S/he felt so tired nothing. | d s/he just sat around and did | | | | | | |
| 4. S/he was very re | estless. | | | | | | |
| 5. S/he felt s/he wa | as no good any more. | | | | | | |
| 6. S/he cried a lot. | | | | | | | |
| 7. S/he found it ha concentrate. | rd to think properly or | | | | | | |
| 8. S/he hated him/ | herself. | | | | | | |
| 9. S/he felt s/he wa | as a bad person. | | | | | | |
| 10. S/he felt lonely. | | | | | | | |

Copyright Adrian Angold & Elizabeth J. Costello, 1987; Developmental Epidemiology Program; Duke University

11. S/he thought nobody really loved him/her.

12. S/he thought s/he could never be as good as

13. S/he felt s/he did everything wrong.

other kids.

Response Scale for SMFQ

0 1 2
Not True Sometimes True