



# TF-CBT Follow Up Forms (Monthly, Periodic, & Discharge) English

Required Forms
1. TF-CBT Monthly Session Form $\square$
2. Child's Behavior & Functioning*  Ohio- Caregiver Report (child 5+) □
3. Child's Trauma Symtoms* $YCPC$ - Caregiver Report $\square$
<b>Note:</b> Alternate or additional measures can be used based on clinical judgment of primary symptom area targeted by treatment
4. Satisfaction Questionnaire (caregiver or child) $\square$
5. Client Discharge Face Sheet $\square$
*Required at periodic and discharge





#### **TF-CBT Monthly Session Form**

#### VALIDATION REQUIREMENTS AND SYMBOLS EXPLAINED

! This symbol means the field is one of the minimum fields that must be filled out to save the record. No data will be saved unless these fields are completed.

This symbol means the field is a required field in order to save the record as completed. Although you can save the record, the system does not consider the record to be completed unless ALL of these fields are completed.

* record to be completed unlo Data Entry Person: Greyed				Client F	ace Sheet-Intake, so y	ou won	't have t	o enter them	again here	
			Direct Service Pro	ovide	r User Informa	ation				
Clinician First Name:				Clini	ician Last Name:					
Project Name:										
			Child	Infor	mation					
First Initial of First Name:			First Initial of Last Name:		1		ate of	Birth:		
			Child Iden	tifica	ation Codes					
Provider Client ID:				PSD	CRS ID:					
			Sessior	ı Info	ormation					
Was there a visit this more (Select one)	/as there a visit this month? delect one)			Yes				No		
<b>Treatment Components</b>										
			Using Measures (administer or share results)		Relaxation			Trauma Narrative Completed		
Please check all Compone Used this month:	ents	with	Case Management (assist with basic needs, collateral contacts with school/DCF, etc.)		Affective Expression		0	In Vivo Exposure		
		Psyc	noeducation		Cognitive Coping			-	ession (prepping or sharing ratives w/caregiver)	
		<b>J</b> Pare	nting Skills		Trauma Narrative			Enhancing	Safety	
Collaboration				1	T			T		
During this month, did yo communicate with the	u 🗀	<b>D</b> CF	Worker		Probation officer			Physician		
child's:		Scho	ol	Other						
Collaboration Notes:		·					•			





#### **TF-CBT Monthly Session Form**

	Functioning								
		Very much improved since the initiation of treatment		Much Improved		Minimally improved			
Compared to the child's condition at the start of TF-CBT, this child's condition is:		No change from baseline (the initiation of treatment)	_	Minimally worse	_	Much Worse			
cor, this child s condition is.		Very much worse since the initiation of treatment							
		Session Fi	delit	y Checklist					
Session Structure									
Prior to how many sessions		None (0%)		Some (34-66%)		All (100%)			
this month did you prepare materials or a session plan?		A few (1-33%)		Most (67-99%)					
During how many sessions		None (0%)		Some (34-66%)		All (100%)			
this month was homework assigned or reviewed?		A few (1-33%)		Most (67-99%)					
During how many sessions		None (0%)		Some (34-66%)		All (100%)			
this month were COWS saved for the end of the session?		A few (1-33%)		Most (67-99%)					
During how many sessions		None (0%)		Some (34-66%)		All (100%)			
this month did the child and/or caregiver practice/demonstrate skill(s) in session (behavior rehearsal)?		A few (1-33%)		Most (67-99%)					

12/5/2019

Client Initials:	Client ID:	Date of Completion:	/	/

## Ohio Mental Health Consumer Outcomes System Ohio Youth Problem and Functioning Scales (Caregiver: English) Parent Rating – Short Form

Instructions: Please rate the degree to which your child has experienced the following problems in the past 30 days.	Not at All	Once or Twice	Several Times	Often	Most of the Time	All of the Time
Arguing with others	0	1	2	3	4	5
2. Getting into fights	0	1	2	3	4	5
3. Yelling, swearing, or screaming at others	0	1	2	3	4	5
4. Fits of anger	0	1	2	3	4	5
5. Refusing to do things teachers or parents ask	0	1	2	3	4	5
6. Causing trouble for no reason	0	1	2	3	4	5
7. Using drugs or alcohol	0	1	2	3	4	5
Breaking rules or breaking the law (out past curfew, stealing)	0	1	2	3	4	5
9. Skipping school or classes	0	1	2	3	4	5
10. Lying	0	1	2	3	4	5
11. Can't seem to sit still, having too much energy	0	1	2	3	4	5
12. Hurting self (cutting or scratching self, taking pills)	0	1	2	3	4	5
13. Talking or thinking about death	0	1	2	3	4	5
14. Feeling worthless or useless	0	1	2	3	4	5
15. Feeling lonely and having no friends	0	1	2	3	4	5
16. Feeling anxious or fearful	0	1	2	3	4	5
17. Worrying that something bad is going to happen	0	1	2	3	4	5
18. Feeling sad or depressed	0	1	2	3	4	5
19. Nightmares	0	1	2	3	4	5
20. Eating problems	0	1	2	3	4	5

Copyright © Benjamin M. Ogles

(Add	ratings	together	) Total	
(2 202 02 .			,	

January 2000 (Parent-1)

## Response Scale for OHIO Problem Scale

0 1 2 3 4 5

Not at all Once or twice times Often Most of the time

Client Initials:	Client ID:	Date of Completion: / /	

#### Ohio Youth Problem and Functioning Scales (Caregiver: English)

Parent Rating – Short Form continued

Instructions: Please rate the degree to which your child's problems affect his or her current ability in everyday activities. Consider your child's current level of functioning.	Extreme Troubles	Quite a Few Troubles	Some Troubles	OK	Doing Very Well
Getting along with friends	0	1	2	3	4
2. Getting along with family	0	1	2	3	4
Dating or developing relationships with boyfriends orgirlfriends	0	1	2	3	4
Getting along with adults outside the family (teachers, principal)	0	1	2	3	4
5. Keeping neat and clean, looking good	0	1	2	3	4
6. Caring for health needs and keeping good health habits (taking medicines or brushing teeth)	0	1	2	3	4
7. Controlling emotions and staying out of trouble	0	1	2	3	4
Being motivated and finishing projects	0	1	2	3	4
Participating in hobbies (baseball cards, coins, stamps, art)	0	1	2	3	4
10. Participating in recreational activities (sports, swimming, bike riding)	0	1	2	3	4
11. Completing household chores (cleaning room, other chores)	0	1	2	3	4
12. Attending school and getting passing grades in school	0	1	2	3	4
13. Learning skills that will be useful for future jobs	0	1	2	3	4
14. Feeling good about self	0	1	2	3	4
15. Thinking clearly and making good decisions	0	1	2	3	4
16. Concentrating, paying attention, and completing tasks	0	1	2	3	4
17. Earning money and learning how to use money wisely	0	1	2	3	4
18. Doing things without supervision or restrictions	0	1	2	3	4
19. Accepting responsibility for actions	0	1	2	3	4
20. Ability to express feelings	0	1	2	3	4

Copyright	©	Benjamin	M.	Ogles
-----------	---	----------	----	-------

January 2000 (Parent-2)

(Add ratings together	r) Total

### Response Scale for OHIO Functioning Scale

0 1 2 3 4

Extreme Quite a few Some OK Doing troubles troubles troubles very well

Client Initials:	Client ID:	Date of Completion: /	/

#### YOUNG CHILD PTSD CHECKLIST (YCPC) (Caregiver: English)

For Child under 7 years old.

Below is a list of symptoms that children can have after life-threatening events.

When you think of ALL the life-threatening traumatic events from the first page, circle the number below (0-4) that best describes how often the symptom has bothered you in the LAST 2 WEEKS.

Not at all   Once in a while   2 to 4 times a week/   Malmost always   Everyday		0	1	2	3			4			
1.       Does your child have intrusive memories of the trauma? Does s/he bring it up on his/her own?       0       1       2       3       4         2.       Does your child re-enact the trauma in play with dolls or toys? This would be seenes that look just like the trauma. Or does s/he act it out by him/herself or with other kids?       0       1       2       3       4         3.       Is your child having more nightmares since the trauma(s) occurred?       0       1       2       3       4         4.       Does your child act like the traumatic event is happening to him/her again, even when it isn't? This is where a child is acting like they are back in the traumatic event and aren't in touch with reality. This is a pretty obvious thing when it happens.       0       1       2       3       4         5.       Since the trauma(s) has s/he had episodes when s/he seems to freeze? You may have tried to snap him/her out of it but s/he was unresponsive.       0       1       2       3       4         6.       Does s/he get upset when exposed to reminders of the event(s)? For example, a child who was in a car wreck might be nervous when it is raining. Or, a child who saw domestic violence might be nervous when it is raining. Or, a child who saw domestic violence might be nervous when other people argue. Or, a girl who was sexually abused might be nervous when other people alk about what happened, does s/he walk away or change the topic?       0       1       2       3       4         8.       Does your c		Not at all		2 to 4 times a week/		eek/		Ever	yday		
Up on his/her own?   2. Does your child re-enact the trauma in play with dolls or toys? This would be scenes that look just like the trauma. Or does s/he act it out by him/herself or with other kids?   3. Is your child having more nightmares since the trauma(s) occurred?   0			Once in a while	Half the time	Almost always						
Up on his/her own?   2. Does your child re-enact the trauma in play with dolls or toys? This would be scenes that look just like the trauma. Or does s/he act it out by him/herself or with other kids?   3. Is your child having more nightmares since the trauma(s) occurred?   0		T =									
Does your child re-enact the trauma in play with dolls or toys? This would be scenes that look just like the trauma. Or does s/he act it out by him/herself or with other kids?   3. Is your child having more nightmares since the trauma(s) occurred?   0   1   2   3   4	1.	-		nories of the trauma?	Does s/he bring it	0	1	2	3	4	
be scenes that look just like the trauma. Or does s/he act it out by him/herself or with other kids?  3. Is your child having more nightmares since the trauma(s) occurred?  4. Does your child act like the traumatic event is happening to him/her again, even when it isn't? This is where a child is acting like they are back in the traumatic event and aren't in touch with reality. This is a pretty obvious thing when it happens.  5. Since the trauma(s) has s/he had episodes when s/he seems to freeze? You may have tried to snap him/her out of it but s/he was unresponsive.  6. Does s/he get upset when exposed to reminders of the event(s)? For example, a child who was in a car wreck might be nervous when it is raining. Or, a child who was in a hurricane might be nervous when other people argue. Or, a girl who was sexually abused might be nervous when someone touches her.  7. Does your child get physically distressed when exposed to reminders? Like heart racing, shaking hands, sweaty, short of breath, or sick to his/her stomach? Think of the same type of examples as in #6.  8. Does your child try to avoid conversations that might remind him/her of the trauma(s)? For example, if other people talk about what happened, does s/he walk away or change the topic?  9. Does your child try to avoid things or places that remind him/her of the trauma(s)? For example, a child who was in a car wreck might try to avoid getting into a car. Or, a child who was in a flood might tell you not to drive over a bridge. Or, a child who was in a flood might tell you not to drive over a bridge. Or, a child who was in a flood might tell you not to drive over a bridge. Or, a child who saw domestic violence might be nervous to go in the house where it occurred. Or, a girl who was sexually abused might be nervous about going to bed because that's where she was abused before.  10. Does your child have difficulty remembering the whole incident? Has s/he blocked out the entire event?  11. Has s/he lost interest in doing things that s/he used to like to		1 1			0.001	-					
him/herself or with other kids?   Is your child having more nightmares since the trauma(s) occurred?   0	2.	_			=	0	1	2	3	4	
<ul> <li>3. Is your child having more nightmares since the trauma(s) occurred?  4. Does your child act like the traumatic event is happening to him/her again, even when it isn't? This is where a child is acting like they are back in the traumatic event and aren't in touch with reality. This is a pretty obvious thing when it happens.</li> <li>5. Since the trauma(s) has s/he had episodes when s/he seems to freeze? You may have tried to snap him/her out of it but s/he was unresponsive.</li> <li>6. Does s/he get upset when exposed to reminders of the event(s)? For example, a child who was in a car wreck might be nervous while riding in a car now. Or, a child who saw domestic violence might be nervous when other people argue. Or, a girl who was sexually abused might be nervous when other people argue. Or, a girl who was sexually abused might be nervous when other people argue. Or, a child who saw domestic violence might be nervous when other people argue. Or, a girl who was sexually abused might be nervous when other people argue. Or, a child who saw domestic violence might be nervous when other people argue. Or, a child voice of breath, or sick to his/her stomach? Think of the same type of examples as in #6.</li> <li>8. Does your child try to avoid conversations that might remind him/her of the trauma(s)? For example, if other people talk about what happened, does s/he walk away or change the topic?</li> <li>9. Does your child try to avoid things or places that remind him/her of the trauma(s)? For example, a child who was in a car wreck might try to avoid getting into a car. Or, a child who was in a flood might tell you not to drive over a bridge. Or, a child who saw domestic violence might be nervous to go in the house where it occurred. Or, a girl who was sexually abused might be nervous about going to bed because that's where she was abused before.</li> <li>10. Does your child have difficulty remembering the whole incident? Has s/he blocked out the entire event?</li> <li>11. Has s/he lost interest in doing thing</li></ul>				ıma. Or does s/he act	it out by						
<ul> <li>4. Does your child act like the traumatic event is happening to him/her again, even when it isn't? This is where a child is acting like they are back in the traumatic event and aren't in touch with reality. This is a pretty obvious thing when it happens.</li> <li>5. Since the trauma(s) has s/he had episodes when s/he seems to freeze? You may have tried to snap him/her out of it but s/he was unresponsive.</li> <li>6. Does s/he get upset when exposed to reminders of the event(s)? For example, a child who was in a car wreck might be nervous while riding in a car now. Or, a child who was in a hurricane might be nervous when other people argue. Or, a girl who was sexually abused might be nervous when someone touches her.</li> <li>7. Does your child get physically distressed when exposed to reminders? Like heart racing, shaking hands, sweaty, short of breath, or sick to his/her stomach? Think of the same type of examples as in #6.</li> <li>8. Does your child try to avoid conversations that might remind him/her of the trauma(s)? For example, if other people talk about what happened, does s/he walk away or change the topic?</li> <li>9. Does your child try to avoid things or places that remind him/her of the trauma(s)? For example, a child who was in a flood might tell you not to drive over a bridge. Or, a child who saw domestic violence might be nervous to go in the house where it occurred. Or, a girl who was sexually abused might be nervous about going to bed because that's where she was abused before.</li> <li>10. Does your child have difficulty remembering the whole incident? Has s/he blocked out the entire event?</li> <li>11. Has s/he lost interest in doing things that s/he used to like to do since the trauma(s)?</li> <li>12. Since the trauma(s), does your child show a restricted range of positive</li> <li>0 1 2 3 4</li> </ul>											
even when it isn't? This is where a child is acting like they are back in the traumatic event and aren't in touch with reality. This is a pretty obvious thing when it happens.  5. Since the trauma(s) has s/he had episodes when s/he seems to freeze? You may have tried to snap him/her out of it but s/he was unresponsive.  6. Does s/he get upset when exposed to reminders of the event(s)? For example, a child who was in a car wreck might be nervous while riding in a car now. Or, a child who saw domestic violence might be nervous when other people argue. Or, a girl who was sexually abused might be nervous when someone touches her.  7. Does your child get physically distressed when exposed to reminders? Like heart racing, shaking hands, sweaty, short of breath, or sick to his/her stomach? Think of the same type of examples as in #6.  8. Does your child try to avoid conversations that might remind him/her of the trauma(s)? For example, if other people talk about what happened, does s/he walk away or change the topic?  9. Does your child try to avoid things or places that remind him/her of the trauma(s)? For example, a child who was in a car wreck might try to avoid getting into a car. Or, a child who was in a flood might tell you not to drive over a bridge. Or, a child who saw domestic violence might be nervous to go in the house where it occurred. Or, a girl who was sexually abused might be nervous about going to bed because that's where she was abused before.  10. Does your child have difficulty remembering the whole incident? Has s/he blocked out the entire event?  11. Has s/he lost interest in doing things that s/he used to like to do since the trauma(s)?  12. Since the trauma(s), does your child show a restricted range of positive  0 1 2 3 4							1				
traumatic event and aren't in touch with reality. This is a pretty obvious thing when it happens.  5. Since the trauma(s) has s/he had episodes when s/he seems to freeze? You may have tried to snap him/her out of it but s/he was unresponsive.  6. Does s/he get upset when exposed to reminders of the event(s)? For example, a child who was in a car wreck might be nervous while riding in a car now. Or, a child who saw domestic violence might be nervous when other people argue. Or, a girl who was sexually abused might be nervous when someone touches her.  7. Does your child get physically distressed when exposed to reminders? Like heart racing, shaking hands, sweaty, short of breath, or sick to his/her stomach? Think of the same type of examples as in #6.  8. Does your child try to avoid conversations that might remind him/her of the trauma(s)? For example, if other people talk about what happened, does s/he walk away or change the topic?  9. Does your child try to avoid things or places that remind him/her of the trauma(s)? For example, a child who was in a car wreck might try to avoid getting into a car. Or, a child who was in a flood might tell you not to drive over a bridge. Or, a child who saw domestic violence might be nervous to go in the house where it occurred. Or, a girl who was sexually abused might be nervous about going to bed because that's where she was abused before.  10. Does your child have difficulty remembering the whole incident? Has s/he blocked out the entire event?  11. Has s/he lost interest in doing things that s/he used to like to do since the trauma(s)?  12. Since the trauma(s), does your child show a restricted range of positive  0 1 2 3 4	4.					0	1	2	3	4	
thing when it happens.  5. Since the trauma(s) has s/he had episodes when s/he seems to freeze? You may have tried to snap him/her out of it but s/he was unresponsive.  6. Does s/he get upset when exposed to reminders of the event(s)? For example, a child who was in a car wreck might be nervous while riding in a car now. Or, a child who was in a hurricane might be nervous when it is raining. Or, a child who saw domestic violence might be nervous when other people argue. Or, a girl who was sexually abused might be nervous when someone touches her.  7. Does your child get physically distressed when exposed to reminders? Like heart racing, shaking hands, sweaty, short of breath, or sick to his/her stomach? Think of the same type of examples as in #6.  8. Does your child try to avoid conversations that might remind him/her of the trauma(s)? For example, if other people talk about what happened, does s/he walk away or change the topic?  9. Does your child try to avoid things or places that remind him/her of the walk away or change the topic?  9. Does your child try to avoid things or places that remind him/her of the trauma(s)? For example, a child who was in a car wreck might try to avoid getting into a car. Or, a child who was in a flood might tell you not to drive over a bridge. Or, a child who was domestic violence might be nervous to go in the house where it occurred. Or, a girl who was sexually abused might be nervous about going to bed because that's where she was abused before.  10. Does your child have difficulty remembering the whole incident? Has s/he blocked out the entire event?  11. Has s/he lost interest in doing things that s/he used to like to do since the trauma(s)?  12. Since the trauma(s), does your child show a restricted range of positive 0 1 2 3 4					-						
5. Since the trauma(s) has s/he had episodes when s/he seems to freeze? You may have tried to snap him/her out of it but s/he was unresponsive.  6. Does s/he get upset when exposed to reminders of the event(s)? For example, a child who was in a car wreck might be nervous while riding in a car now. Or, a child who was in a hurricane might be nervous when it is raining. Or, a child who saw domestic violence might be nervous when other people argue. Or, a girl who was sexually abused might be nervous when someone touches her.  7. Does your child get physically distressed when exposed to reminders? Like heart racing, shaking hands, sweaty, short of breath, or sick to his/her stomach? Think of the same type of examples as in #6.  8. Does your child try to avoid conversations that might remind him/her of the trauma(s)? For example, if other people talk about what happened, does s/he walk away or change the topic?  9. Does your child try to avoid things or places that remind him/her of the value and thing into a car. Or, a child who was in a car wreck might try to avoid getting into a car. Or, a child who was in a flood might tell you not to drive over a bridge. Or, a child who saw domestic violence might be nervous to go in the house where it occurred. Or, a girl who was sexually abused might be nervous about going to bed because that's where she was abused before.  10. Does your child have difficulty remembering the whole incident? Has s/he blocked out the entire event?  11. Has s/he lost interest in doing things that s/he used to like to do since the trauma(s)?  12. Since the trauma(s), does your child show a restricted range of positive  0 1 2 3 4				with reality. This is a	pretty obvious						
may have tried to snap him/her out of it but s/he was unresponsive.  6. Does s/he get upset when exposed to reminders of the event(s)? For example, a child who was in a car wreck might be nervous while riding in a car now. Or, a child who was in a hurricane might be nervous when it is raining. Or, a child who saw domestic violence might be nervous when other people argue. Or, a girl who was sexually abused might be nervous when someone touches her.  7. Does your child get physically distressed when exposed to reminders? Like heart racing, shaking hands, sweaty, short of breath, or sick to his/her stomach? Think of the same type of examples as in #6.  8. Does your child try to avoid conversations that might remind him/her of the trauma(s)? For example, if other people talk about what happened, does s/he walk away or change the topic?  9. Does your child try to avoid things or places that remind him/her of the trauma(s)? For example, a child who was in a car wreck might try to avoid getting into a car. Or, a child who was in a flood might tell you not to drive over a bridge. Or, a child who saw domestic violence might be nervous to go in the house where it occurred. Or, a girl who was sexually abused might be nervous about going to bed because that's where she was abused before.  10. Does your child have difficulty remembering the whole incident? Has s/he blocked out the entire event?  11. Has s/he lost interest in doing things that s/he used to like to do since the trauma(s)?  12. Since the trauma(s), does your child show a restricted range of positive  0 1 2 3 4											
6. Does s/he get upset when exposed to reminders of the event(s)? For example, a child who was in a car wreck might be nervous while riding in a car now. Or, a child who saw domestic violence might be nervous when other people argue. Or, a girl who was sexually abused might be nervous when someone touches her.  7. Does your child get physically distressed when exposed to reminders? Like heart racing, shaking hands, sweaty, short of breath, or sick to his/her stomach? Think of the same type of examples as in #6.  8. Does your child try to avoid conversations that might remind him/her of the trauma(s)? For example, if other people talk about what happened, does s/he walk away or change the topic?  9. Does your child try to avoid things or places that remind him/her of the trauma(s)? For example, a child who was in a car wreck might try to avoid getting into a car. Or, a child who was in a flood might tell you not to drive over a bridge. Or, a child who saw domestic violence might be nervous to go in the house where it occurred. Or, a girl who was sexually abused might be nervous about going to bed because that's where she was abused before.  10. Does your child have difficulty remembering the whole incident? Has s/he blocked out the entire event?  11. Has s/he lost interest in doing things that s/he used to like to do since the trauma(s)?  12. Since the trauma(s), does your child show a restricted range of positive  0 1 2 3 4	5.					0	1	2	3	4	
example, a child who was in a car wreck might be nervous while riding in a car now. Or, a child who was in a hurricane might be nervous when it is raining. Or, a child who saw domestic violence might be nervous when other people argue. Or, a girl who was sexually abused might be nervous when someone touches her.  7. Does your child get physically distressed when exposed to reminders? Like heart racing, shaking hands, sweaty, short of breath, or sick to his/her stomach? Think of the same type of examples as in #6.  8. Does your child try to avoid conversations that might remind him/her of the trauma(s)? For example, if other people talk about what happened, does s/he walk away or change the topic?  9. Does your child try to avoid things or places that remind him/her of the trauma(s)? For example, a child who was in a car wreck might try to avoid getting into a car. Or, a child who was in a flood might tell you not to drive over a bridge. Or, a child who saw domestic violence might be nervous to go in the house where it occurred. Or, a girl who was sexually abused might be nervous about going to bed because that's where she was abused before.  10. Does your child have difficulty remembering the whole incident? Has s/he blocked out the entire event?  11. Has s/he lost interest in doing things that s/he used to like to do since the trauma(s)?											
car now. Or, a child who was in a hurricane might be nervous when it is raining. Or, a child who saw domestic violence might be nervous when other people argue. Or, a girl who was sexually abused might be nervous when someone touches her.  7. Does your child get physically distressed when exposed to reminders? Like heart racing, shaking hands, sweaty, short of breath, or sick to his/her stomach? Think of the same type of examples as in #6.  8. Does your child try to avoid conversations that might remind him/her of the trauma(s)? For example, if other people talk about what happened, does s/he walk away or change the topic?  9. Does your child try to avoid things or places that remind him/her of the trauma(s)? For example, a child who was in a car wreck might try to avoid getting into a car. Or, a child who was in a flood might tell you not to drive over a bridge. Or, a child who saw domestic violence might be nervous to go in the house where it occurred. Or, a girl who was sexually abused might be nervous about going to bed because that's where she was abused before.  10. Does your child have difficulty remembering the whole incident? Has s/he blocked out the entire event?  11. Has s/he lost interest in doing things that s/he used to like to do since the trauma(s)?  12. Since the trauma(s), does your child show a restricted range of positive  0 1 2 3 4	6.					0	1	2	3	4	
raining. Or, a child who saw domestic violence might be nervous when other people argue. Or, a girl who was sexually abused might be nervous when someone touches her.  7. Does your child get physically distressed when exposed to reminders? Like heart racing, shaking hands, sweaty, short of breath, or sick to his/her stomach? Think of the same type of examples as in #6.  8. Does your child try to avoid conversations that might remind him/her of the trauma(s)? For example, if other people talk about what happened, does s/he walk away or change the topic?  9. Does your child try to avoid things or places that remind him/her of the trauma(s)? For example, a child who was in a car wreck might try to avoid getting into a car. Or, a child who was in a flood might tell you not to drive over a bridge. Or, a child who saw domestic violence might be nervous to go in the house where it occurred. Or, a girl who was sexually abused might be nervous about going to bed because that's where she was abused before.  10. Does your child have difficulty remembering the whole incident? Has s/he blocked out the entire event?  11. Has s/he lost interest in doing things that s/he used to like to do since the trauma(s)?  12. Since the trauma(s), does your child show a restricted range of positive  0 1 2 3 4		example, a ch	ild who was in a car v	wreck might be nervou	is while riding in a						
other people argue. Or, a girl who was sexually abused might be nervous when someone touches her.  7. Does your child get physically distressed when exposed to reminders? Like heart racing, shaking hands, sweaty, short of breath, or sick to his/her stomach? Think of the same type of examples as in #6.  8. Does your child try to avoid conversations that might remind him/her of the trauma(s)? For example, if other people talk about what happened, does s/he walk away or change the topic?  9. Does your child try to avoid things or places that remind him/her of the trauma(s)? For example, a child who was in a car wreck might try to avoid getting into a car. Or, a child who was in a flood might tell you not to drive over a bridge. Or, a child who saw domestic violence might be nervous to go in the house where it occurred. Or, a girl who was sexually abused might be nervous about going to bed because that's where she was abused before.  10. Does your child have difficulty remembering the whole incident? Has s/he blocked out the entire event?  11. Has s/he lost interest in doing things that s/he used to like to do since the trauma(s)?  12. Since the trauma(s), does your child show a restricted range of positive  0 1 2 3 4				•							
when someone touches her.  7. Does your child get physically distressed when exposed to reminders? Like heart racing, shaking hands, sweaty, short of breath, or sick to his/her stomach? Think of the same type of examples as in #6.  8. Does your child try to avoid conversations that might remind him/her of the trauma(s)? For example, if other people talk about what happened, does s/he walk away or change the topic?  9. Does your child try to avoid things or places that remind him/her of the trauma(s)? For example, a child who was in a car wreck might try to avoid getting into a car. Or, a child who was in a flood might tell you not to drive over a bridge. Or, a child who saw domestic violence might be nervous to go in the house where it occurred. Or, a girl who was sexually abused might be nervous about going to bed because that's where she was abused before.  10. Does your child have difficulty remembering the whole incident? Has s/he blocked out the entire event?  11. Has s/he lost interest in doing things that s/he used to like to do since the trauma(s)?  12. Since the trauma(s), does your child show a restricted range of positive  0 1 2 3 4		raining. Or, a	child who saw domes	tic violence might be	nervous when						
<ul> <li>7. Does your child get physically distressed when exposed to reminders? Like heart racing, shaking hands, sweaty, short of breath, or sick to his/her stomach? Think of the same type of examples as in #6.</li> <li>8. Does your child try to avoid conversations that might remind him/her of the trauma(s)? For example, if other people talk about what happened, does s/he walk away or change the topic?</li> <li>9. Does your child try to avoid things or places that remind him/her of the trauma(s)? For example, a child who was in a car wreck might try to avoid getting into a car. Or, a child who was in a flood might tell you not to drive over a bridge. Or, a child who saw domestic violence might be nervous to go in the house where it occurred. Or, a girl who was sexually abused might be nervous about going to bed because that's where she was abused before.</li> <li>10. Does your child have difficulty remembering the whole incident? Has s/he blocked out the entire event?</li> <li>11. Has s/he lost interest in doing things that s/he used to like to do since the trauma(s)?</li> <li>12. Since the trauma(s), does your child show a restricted range of positive</li> <li>0 1 2 3 4</li> </ul>				vas sexually abused m	night be nervous						
heart racing, shaking hands, sweaty, short of breath, or sick to his/her stomach? Think of the same type of examples as in #6.  8. Does your child try to avoid conversations that might remind him/her of the trauma(s)? For example, if other people talk about what happened, does s/he walk away or change the topic?  9. Does your child try to avoid things or places that remind him/her of the trauma(s)? For example, a child who was in a car wreck might try to avoid getting into a car. Or, a child who was in a flood might tell you not to drive over a bridge. Or, a child who saw domestic violence might be nervous to go in the house where it occurred. Or, a girl who was sexually abused might be nervous about going to bed because that's where she was abused before.  10. Does your child have difficulty remembering the whole incident? Has s/he blocked out the entire event?  11. Has s/he lost interest in doing things that s/he used to like to do since the trauma(s)?  12. Since the trauma(s), does your child show a restricted range of positive  0 1 2 3 4		when someon	e touches her.								
stomach? Think of the same type of examples as in #6.  8. Does your child try to avoid conversations that might remind him/her of the trauma(s)? For example, if other people talk about what happened, does s/he walk away or change the topic?  9. Does your child try to avoid things or places that remind him/her of the trauma(s)? For example, a child who was in a car wreck might try to avoid getting into a car. Or, a child who was in a flood might tell you not to drive over a bridge. Or, a child who saw domestic violence might be nervous to go in the house where it occurred. Or, a girl who was sexually abused might be nervous about going to bed because that's where she was abused before.  10. Does your child have difficulty remembering the whole incident? Has s/he blocked out the entire event?  11. Has s/he lost interest in doing things that s/he used to like to do since the trauma(s)?  12. Since the trauma(s), does your child show a restricted range of positive  0 1 2 3 4	7.	Does your chi	ild get physically distr	essed when exposed t	o reminders? Like	0	1	2	3	4	
8. Does your child try to avoid conversations that might remind him/her of the trauma(s)? For example, if other people talk about what happened, does s/he walk away or change the topic?  9. Does your child try to avoid things or places that remind him/her of the trauma(s)? For example, a child who was in a car wreck might try to avoid getting into a car. Or, a child who was in a flood might tell you not to drive over a bridge. Or, a child who saw domestic violence might be nervous to go in the house where it occurred. Or, a girl who was sexually abused might be nervous about going to bed because that's where she was abused before.  10. Does your child have difficulty remembering the whole incident? Has s/he blocked out the entire event?  11. Has s/he lost interest in doing things that s/he used to like to do since the trauma(s)?  12. Since the trauma(s), does your child show a restricted range of positive  0 1 2 3 4		heart racing, s	shaking hands, sweaty	, short of breath, or si	ck to his/her						
trauma(s)? For example, if other people talk about what happened, does s/he walk away or change the topic?  9. Does your child try to avoid things or places that remind him/her of the trauma(s)? For example, a child who was in a car wreck might try to avoid getting into a car. Or, a child who was in a flood might tell you not to drive over a bridge. Or, a child who saw domestic violence might be nervous to go in the house where it occurred. Or, a girl who was sexually abused might be nervous about going to bed because that's where she was abused before.  10. Does your child have difficulty remembering the whole incident? Has s/he blocked out the entire event?  11. Has s/he lost interest in doing things that s/he used to like to do since the trauma(s)?  12. Since the trauma(s), does your child show a restricted range of positive  0 1 2 3 4											
walk away or change the topic?  9. Does your child try to avoid things or places that remind him/her of the trauma(s)? For example, a child who was in a car wreck might try to avoid getting into a car. Or, a child who was in a flood might tell you not to drive over a bridge. Or, a child who saw domestic violence might be nervous to go in the house where it occurred. Or, a girl who was sexually abused might be nervous about going to bed because that's where she was abused before.  10. Does your child have difficulty remembering the whole incident? Has s/he blocked out the entire event?  11. Has s/he lost interest in doing things that s/he used to like to do since the trauma(s)?  12. Since the trauma(s), does your child show a restricted range of positive  0 1 2 3 4	8.	Does your chi	ild try to avoid conver	sations that might ren	nind him/her of the	0	1	2	3	4	
9. Does your child try to avoid things or places that remind him/her of the trauma(s)? For example, a child who was in a car wreck might try to avoid getting into a car. Or, a child who was in a flood might tell you not to drive over a bridge. Or, a child who saw domestic violence might be nervous to go in the house where it occurred. Or, a girl who was sexually abused might be nervous about going to bed because that's where she was abused before.  10. Does your child have difficulty remembering the whole incident? Has s/he blocked out the entire event?  11. Has s/he lost interest in doing things that s/he used to like to do since the trauma(s)?  12. Since the trauma(s), does your child show a restricted range of positive  0 1 2 3 4		trauma(s)? Fo	or example, if other pe	ople talk about what h	nappened, does s/he						
trauma(s)? For example, a child who was in a car wreck might try to avoid getting into a car. Or, a child who was in a flood might tell you not to drive over a bridge. Or, a child who saw domestic violence might be nervous to go in the house where it occurred. Or, a girl who was sexually abused might be nervous about going to bed because that's where she was abused before.  10. Does your child have difficulty remembering the whole incident? Has s/he blocked out the entire event?  11. Has s/he lost interest in doing things that s/he used to like to do since the trauma(s)?  12. Since the trauma(s), does your child show a restricted range of positive  0 1 2 3 4		walk away or	change the topic?								
getting into a car. Or, a child who was in a flood might tell you not to drive over a bridge. Or, a child who saw domestic violence might be nervous to go in the house where it occurred. Or, a girl who was sexually abused might be nervous about going to bed because that's where she was abused before.  10. Does your child have difficulty remembering the whole incident? Has s/he blocked out the entire event?  11. Has s/he lost interest in doing things that s/he used to like to do since the trauma(s)?  12. Since the trauma(s), does your child show a restricted range of positive  0 1 2 3 4	9.	Does your chi	ild try to avoid things	or places that remind	him/her of the	0	1	2	3	4	
over a bridge. Or, a child who saw domestic violence might be nervous to go in the house where it occurred. Or, a girl who was sexually abused might be nervous about going to bed because that's where she was abused before.  10. Does your child have difficulty remembering the whole incident? Has s/he blocked out the entire event?  11. Has s/he lost interest in doing things that s/he used to like to do since the trauma(s)?  12. Since the trauma(s), does your child show a restricted range of positive  0 1 2 3 4		trauma(s)? Fo	or example, a child wh	o was in a car wreck	might try to avoid						
go in the house where it occurred. Or, a girl who was sexually abused might be nervous about going to bed because that's where she was abused before.  10. Does your child have difficulty remembering the whole incident? Has s/he blocked out the entire event?  11. Has s/he lost interest in doing things that s/he used to like to do since the trauma(s)?  12. Since the trauma(s), does your child show a restricted range of positive  13. 4		getting into a	car. Or, a child who v	vas in a flood might te	ell you not to drive						
be nervous about going to bed because that's where she was abused before.  10. Does your child have difficulty remembering the whole incident? Has s/he blocked out the entire event?  11. Has s/he lost interest in doing things that s/he used to like to do since the trauma(s)?  12. Since the trauma(s), does your child show a restricted range of positive  13. 4		over a bridge.	Or, a child who saw	domestic violence mig	ght be nervous to						
10. Does your child have difficulty remembering the whole incident? Has s/he blocked out the entire event?  11. Has s/he lost interest in doing things that s/he used to like to do since the trauma(s)?  12. Since the trauma(s), does your child show a restricted range of positive  13. 4		go in the hous	se where it occurred. (	Or, a girl who was sex	ually abused might						
blocked out the entire event?  11. Has s/he lost interest in doing things that s/he used to like to do since the trauma(s)?  12. Since the trauma(s), does your child show a restricted range of positive  13. The property of the trauma(s) of the											
11. Has s/he lost interest in doing things that s/he used to like to do since the trauma(s)?  12. Since the trauma(s), does your child show a restricted range of positive  13. 4	10.			nembering the whole i	ncident? Has s/he						
trauma(s)?  12. Since the trauma(s), does your child show a restricted range of positive  0 1 2 3 4		blocked out th	ne entire event?								
12. Since the trauma(s), does your child show a restricted range of positive 0 1 2 3 4	11.	Has s/he lost	interest in doing thing	s that s/he used to like	e to do since the	0	1	2	3	4	
		trauma(s)?									
emotions on his/her face compared to before?	12.	Since the trau	ma(s), does your child	d show a restricted rar	nge of positive	0	1	2	3	4	
		emotions on h	nis/her face compared	to before?							

Client Initials:	Client ID:	Date of Completion: /	/

#### YCPC (Caregiver: English) continued

13.	Has your child lost hope for the future? For example, s/he believes will not					
14.	have fun tomorrow, or will never be good at anything.  Since the trauma(s) has your child become more distant and withdrawn	0	1	2	3	4
14.	from family members, relatives, or friends?	U	1	2	3	4
15.	Has s/he had a hard time falling asleep or staying asleep since the	0	1	2	3	4
	trauma(s)?					
16.	Has your child become more irritable, or had outbursts of anger, or	0	1	2	3	4
	developed extreme temper tantrums since the trauma(s)?					
17.	Has your child had more trouble concentrating since the trauma(s)?	0	1	2	3	4
18.	Has s/he been more "on the alert" for bad things to happen? For example,	0	1	2	3	4
	does s/he look around for danger?					
19.	Does your child startle more easily than before the trauma(s)? For example,	0	1	2	3	4
	if there's a loud noise or someone sneaks up behind him/her, does s/he jump					
	or seem startled?					
20.	Has your child become more physically aggressive since the trauma(s)?	0	1	2	3	4
	Like hitting, kicking, biting, or breaking things.					
21.	Has s/he become more clingy to you since the trauma(s)?	0	1	2	3	4
22.	Did night terrors start or get worse after the trauma(s)? Night terrors are	0	1	2	3	4
	different from nightmares: in night terrors a child usually screams in their					
	sleep, they don't wake up, and they don't remember it the next day.					
23.	Since the trauma(s), has your child lost previously acquired skills? For	0	1	2	3	4
	example, lost toilet training? Or, lost language skills? Or, lost motor skills					
	working snaps, buttons, or zippers?					
24.	Since the trauma(s), has your child developed any new fears about things	0	1	2	3	4
	that <u>don't seem related</u> to the trauma(s)? What about going to the bathroom					
	alone? Or, being afraid of the dark?					
	FUNCTIONAL IMPAIRMENT					
	Do the symptoms that you endorsed above get in the way of your child's					
	ability to function in the following areas?					
25.	Do (symptoms) substantially "get in the way" of how s/he gets along with	0	1	2	3	4
	you, interfere in your relationship, or make you feel upset or annoyed?					
26.	Do these (symptoms) "get in the way" of how s/he gets along with brothers	0	1	2	3	4
	or sisters, and make them feel upset or annoyed?					
27.	Do (symptoms) "get in the way" of how s/he gets along with friends at all –	0	1	2	3	4
	at daycare, school, or in your neighborhood?					
28.	Do these (symptoms) "get in the way" with the teacher or the class more	0	1	2	3	4
	than average?					
29.	Do (symptoms) make it harder for you to take him/her out in public than it	0	1	2	3	4
	would be with an average child?					
	Is it harder to go out with your child to places like the grocery store? Or to a					
	restaurant?				_	
30.	Do you think that these behaviors cause your child to feel upset?	0	1	2	3	4

<sup>©</sup> Michael Scheeringa, MD, MPH, 2010, Tulane University, New Orleans, LA. <u>mscheer@tulane.edu</u>. This form may be reproduced and used for free, but not sold, without further permission from the author.

### Young Child PTSD Checklist Caregiver Response Scale

Client Initials:	Climate		/ /	,
i lient initials.	Client ID:	Date of Completion: /	,	
Circuit initials.	CHCHCID.	bate of completion. /	/	



#### **Satisfaction Questionnaire**

P

#### Parent Rating -OHIO SATISFACTION SCALE (English)

Instructions: Please circle your response to each question.

- 1. Extremely satisfied
- 2. Moderately satisfied
- 3. Somewhat satisfied
- 4. Somewhat dissatisfied
- 5. Moderately dissatisfied
- 6. Extremely dissatisfied

2. To what degree have you been included in the treatment planning process for y	your child?
--	-------------

- 1. A great deal
- 2. Quite a bit
- 3. Moderately
- 4. Somewhat
- 5. A little
- 6. Not at all

3. Mental health v	workers involved in r	ny case listen	to and value m	ny ideas about t	reatment planning
for my child.					

- 1. A great deal
- 2. Quite a bit
- 3. Moderately
- 4. Somewhat
- 5. A little
- 6. Not at all

#### 4. To what extent does your child's treatment plan include your ideas about your child's treatment needs?

- 1. A great deal
- 2. Quite a bit
- 3. Moderately
- 4. Somewhat
- 5. A little
- 6. Not at all

Total:			
ı Otai.			

Stranger to the transport	CP: LID	Data (Carallatia)	,
Client Initials:	Client ID:	Date of Completion: /	,
cheffe fillerais.	CIICITE ID.	Date of completion.	,





Satisfaction Questionnaire	Y
Youth Rating - OHIO SATISFACTION SCALE	
Form Completed By: ☐ Caregiver ☐ Child ☐ Other:	
Instructions: Please circle your response to each question.	
1. How satisfied are you with the mental health services you have received so far?	
<ol> <li>Extremely satisfied</li> <li>Moderately satisfied</li> <li>Somewhat satisfied</li> <li>Somewhat dissatisfied</li> <li>Moderately dissatisfied</li> <li>Extremely dissatisfied</li> </ol>	
2. How much are you included in deciding your treatment?	
<ol> <li>A great deal</li> <li>Quite a bit</li> <li>Moderately</li> <li>Somewhat</li> <li>A little</li> <li>Not at all</li> </ol>	
3. Mental health workers involved in my case listen to me and know what I want.	
<ol> <li>A great deal</li> <li>Quite a bit</li> <li>Moderately</li> <li>Somewhat</li> <li>A little</li> <li>Not at all</li> </ol>	
4. I have a lot of say about what happens in my treatment.	
<ol> <li>A great deal</li> <li>Quite a bit</li> <li>Moderately</li> <li>Somewhat</li> <li>A little</li> </ol>	

6. Not at all





#### **Discharge Facesheet (MATCH-ADTC & TF-CBT)**

! This symbol means the field is one of the minimum fields that must be filled out to save the record. No data will be saved unless these fields are completed.

This symbol means the field is a required field in order to save the record as completed. Although you can save the record, the system does not consider the record to be completed unless ALL of these fields are completed.

Data Entry Person: Greyed-out fields are pulled in from the completed Client Face Sheet-Intake, so you won't have to enter them again here							
		Direc	t Service Provid	er L	Iser Information		
Clinician First Name:				Clin	ician Last Name:		
Project:				Trea	atment Model Site:		
			Child Info	rma	ation		
Grade (current): *							
			Child Identific	atio	on Codes		
Provider's Unique Client ID:				PSD	CRS ID:		
Which EBP?		MATCH-	ADTC		TF-CBT		
	Discharge Information						
How many visits during this case:			Discharge Date: *	·	//		
% of the total time spent with the child ONLY during this case:	The total time spent for these three % questions should equal 100%				should equal 100%		
% of the total time spent with the caregiver ONLY during this case:		The total time spent for these three % questions should equal 100%					should equal 100%
% of the total time spent with the child and caregiver TOGETHER during this case:			The total time spent for these three % questions should equal 100%				should equal 100%
CGI: Considering your experience, how severe are the child's emotional, behavioral, and/or cognitive concerns at Discharge? (Circle one): *	A	Among the	Normal Slightly Severe Mildly Severe Moderately Severe Markedly Severe Very Severe most severe symptoms to	:hat	CGI: Compared to the child's condition at intake, this child's condition is (Circle one): *		Very much improved  Much improved  Minimally improved  No change  Minimally worse  Much worse  Very much worse
		EBP Mode	ly completed selected el requirements-no itment needed		Referred for other EBP (outpatient) within agency	О	Family moved out of area
Discharge Reason: *		EBP Mode	lly completed selected el requirements- with other treatment		Referred for other non-EBP (outpatient) within agency		Referred to other agency (outpatient)
		Family dis	continued treatment		Referred to higher level of care	П	Assessment Only-no treatment needed
	Other (specify):						





#### Discharge Facesheet (MATCH-ADTC & TF-CBT)

System Involvement								
Child/Family involved with DCF? *				Yes		No		
If child / family is involved with DCF, please complete ALL of the following questions:								
DCF Case ID: (if available)	DCF Case ID: (if available)		DCF Person Link ID: (if available)					
		Child Protective Services – In- Home		Family with Service Needs – (FWSN) In-Home		Not DCF – On Probation		
DCF Status:		Child Protective Services – Out of Home		Family with Service Needs (FWSN) Out of Home		Not DCF – Other Court Involved		
DCF Regional Office:		Dual Commitment (JJ and Child Protective Services)		Juvenile Justice (delinquency) commitment		Termination of Parental Rights		
		Family Assessment Response		Not DCF		Voluntary Services Program		
Youth involved with Juvenile Justic	e (11)	System? *		Yes		No		
If youth is involved with JJ, please of	omp	olete ALL of the following qu	uesti	ons:				
CSSD Client ID: (if available)			css	D Case ID: (if available)				
CSSD Case Type:				Delinquency		Family with Service Needs (Status Offense)		
		Administrative Supervision		Juvenile probation		Restore Probation		
CSSD Case Status:		Extended Probation		Non-Judicial FWSN Family Service Agreement		Suspended Order		
		Interim Orders		Non-Judicial Supervision (NJS)		Waived PDS - Probation		
		Judicial FWSN Supervision	Non-Judicial Supervision Agreement					
Court District:								
Court Handling Decision:				Judicial		Non-Judicial		
		Treatment Infor	mat	ion: School				
Since the start of EBP treatment		,						
Child's school attendance: *		Good (few or no days missed)		No School Attendance: Child Too Young for School		No School Attendance: Other		
		Fair (several days missed)		No School Attendance: Child Suspended/Expelled from School				
		Poor (many days missed)		No School Attendance: Child Dropped Out of School				
Suspended or expelled: *				Yes		No		
<b>IEP:</b> *Does the child have an Individual Edu	catio	n Plan (special education)?		Yes		No		
		Treatment Info	rma	tion: Legal				
Since the start of EBP treatment								
Arrested: * Has the child been arrested s	ince s	start of treatment?		Yes		No		
<b>Detained or incarcerated: *</b> Has the since start of treatment?	child	been detained or incarcerated		Yes		No		
Treatment Information: Medical								





#### **Discharge Facesheet (MATCH-ADTC & TF-CBT)**

Since the start of EBP treatment							
Alcohol and/or drugs problems: *		Yes		No			
Evaluated in ER/ED for psychiatric issues: *		Yes		No			
Certified medically complex: *		Yes		No			

Rev 6/30/2020